

Name: SOLUTIONS

Math 141- Midterm Exam #2 - October 24, 2007

1. (50 points) Find $\frac{dy}{dx}$. You do not need to simplify your answers.

a. $y = x \cos(x)$

$$y' = \cos x - x \sin x$$

b. $y = \tan(x^2 + 1)$

$$y' = 2x \sec^2(x^2 + 1)$$

c. $y = \frac{\sqrt{x}}{x^3 + 1}$

$$y' = \frac{(x^3 + 1)\left(\frac{1}{2\sqrt{x}}\right) - \sqrt{x} \cdot 3x^2}{(x^3 + 1)^2}$$

d. $y = \ln(5x - 15)$

$$y' = \frac{5}{5x - 15}$$

e. $y = 3^x$

$$y' = 3^x \ln 3$$

f. $y = x^{10} - 9x^3 + 15x - 3$

$$y' = 10x^9 - 27x^2 + 15$$

g. $y = \sin^3(x^2)$

$$y' = 3\sin^2(x^2)\cos(x^2) \cdot 2x$$

h. $y = (\sin x)^x$

$$\ln y = x \ln \sin x$$

$$\frac{1}{y} y' = \ln \sin x + \frac{x - \cos x}{\sin x}$$

$$y' = (\sin x)^x \cdot (\ln \sin x + x \cot x)$$

i. $y = \sqrt{\frac{(x^2+1)^5 e^x x^9}{x^2+2}}$

$$\ln y = \frac{1}{2} (5 \ln(x^2+1) + x \ln e + 9 \ln x - \ln(x^2+2))$$

$$\frac{1}{y} y' = \frac{1}{2} \left(\frac{10x}{x^2+1} + 1 + \frac{9}{x} - \frac{2x}{x^2+2} \right)$$

$$y' = \frac{1}{2} \sqrt{\frac{(x^2+1)^5 e^x x^9}{x^2+2}} \left(\frac{10x}{x^2+1} + 1 + \frac{9}{x} - \frac{2x}{x^2+2} \right)$$

j. $\sin(xy) = 5x$

$$\cos(xy) (y + xy') = 5$$

$$y \cos(xy) + y' x \cos(xy) = 5$$

$$y' = \frac{5 - y \cos(xy)}{x \cos(xy)}$$

2. (10 points) Find the equation of the tangent line to the curve

$$2x - xy^2 = -6$$

at the point (3, 2).

$$2 - y^2 - x \cdot 2y y' = 0$$

$$2 - 4 - 12y' = 0$$

$$y' = -2/12 = -1/6$$

$$y - 2 = -\frac{1}{6}(x - 3)$$

3. (10 points) Find the linear approximation to the function $f(x) = x^{3/4}$ at $x=16$. Then use this linear approximation to estimate $15^{3/4}$.

$$f(x) = x^{3/4} \quad f'(x) = \frac{3}{4} x^{-1/4} = \frac{3}{4\sqrt[4]{x}}$$

$$f(16) = 16^{3/4} = 8 \quad f'(16) = \frac{3}{4 \cdot 2} = 3/8$$

$$L(x) = \frac{3}{8}(x - 16) + 8$$

$$15^{3/4} = f(15) \approx L(15) = \frac{3}{8}|15 - 16| + 8$$

$$= -3/8 + 8$$

$$= 61/8$$

4. (5 points) Evaluate this limit by first expressing it as a derivative:

$$\lim_{h \rightarrow 0} \frac{\sqrt[4]{16+h} - 2}{h}$$

If $f(x) = x^{1/4}$ then this limit is $f'(16)$

$$f'(x) = \frac{1}{4} x^{-3/4} = \frac{1}{4(\sqrt[4]{x})^3}$$

$$f'(16) = \left(\frac{1}{32} \right)$$

5. (10 points) Let $r(x) = f(g(h(x)))$, where $h(1) = 2$, $h'(2) = -1$, $g(2) = 3$, $h'(1) = 4$, $g'(2) = 5$, and $f'(3) = 6$. Find $r'(1)$.

$$r'(x) = f'(g(h(x))) \cdot g'(h(x)) \cdot h'(x)$$


$$= f'(g(2)) \cdot g'(2) \cdot 4$$

$$= f'(3) \cdot 5 \cdot 4$$

$$= 6 \cdot 5 \cdot 4$$

$$= \left(120 \right)$$

6. (15 points) The angle of elevation of the sun is decreasing at a rate of 0.25 radians/hour. How fast is the length of the shadow cast by a 4 foot tall pole increasing when the angle of elevation of the sun is $\pi/6$? (FYI: $\cos(\pi/6) = \sqrt{3}/2$, $\sin(\pi/6) = 1/2$.)




Given: $\frac{d\theta}{dt} = -0.25$

Find $\frac{ds}{dt}$ when $\theta = \pi/6$

$$\tan \theta = \frac{4}{s}$$

$$\sec^2 \theta \frac{d\theta}{dt} = \frac{-4}{s^2} \frac{ds}{dt}$$

When $\theta = \pi/6$:



$$\tan 30 = 1/\sqrt{3}$$

$$\frac{4}{s} = \frac{1}{\sqrt{3}} \rightarrow s = 4\sqrt{3}$$

$$\sec^2(\pi/6) = \frac{1}{\cos^2(\pi/6)} = \frac{1}{3/4} = 4/3$$

$$\rightarrow \frac{4}{3} \left(-\frac{1}{4}\right) = \frac{-4}{48} \cdot \frac{ds}{dt}$$

$$\frac{1}{3} = \frac{1}{12} \frac{ds}{dt}$$

$$\frac{ds}{dt} = 4 \text{ feet/hour}$$