



Non-Sexist Solution of the Menage Problem

Author(s): Kenneth P. Bogart and Peter G. Doyle

Source: *The American Mathematical Monthly*, Vol. 93, No. 7 (Aug. - Sep., 1986), pp. 514-518

Published by: Mathematical Association of America

Stable URL: <http://www.jstor.org/stable/2323022>

Accessed: 05-12-2016 13:32 UTC

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at

<http://about.jstor.org/terms>



Mathematical Association of America is collaborating with JSTOR to digitize, preserve and extend access to *The American Mathematical Monthly*

proof in their own words, as mathematicians do when they try to understand new material. They also treated the case of equality in the Bieberbach conjecture [8], as would others. In the mean time, de Branges produced a more sophisticated write-up of his proof which includes the case of equality in a very natural way [4]. Comments on the exciting events have been written up by FitzGerald [7] and by Gautschi [9], among others. (Added in proof: see also Pommerenke [15].)

References

1. R. Askey and G. Gasper, Positive Jacobi polynomial sums II, *Amer. J. Math.*, 98 (1976) 709–737.
2. L. Bieberbach, Über die Koeffizienten derjenigen Potenzreihen, welche eine schlichte Abbildung des Einheitskreises vermitteln, *S.-B. Preuss. Akad. Wiss.*, 1916, pp. 940–955.
3. L. de Branges, A proof of the Bieberbach conjecture, preprint E-5-84, Steklov Math. Inst., Leningrad Branch, 1984, pp. 1–21.
4. ———, A proof of the Bieberbach conjecture, *Acta Math.*, 154 (1985) 137–152.
5. ———, Square summable power series, 2nd ed., to appear.
6. P. L. Duren, *Univalent Functions*, Springer, Berlin, 1983.
7. C. H. FitzGerald, The Bieberbach conjecture: retrospective. *Notices, Amer. Math. Soc.*, 32 (1985) 2–6.
8. C. H. FitzGerald and C. Pommerenke, The de Branges theorem on univalent functions, *Trans. Amer. Math. Soc.*, 290 (1985) 683–690.
9. W. Gautschi, Reminiscences of my involvement in de Branges' proof of the Bieberbach conjecture, to appear.
10. G. M. Goluzin, *Geometric theory of functions of a complex variable*. English transl., *Amer. Math. Soc.*, Providence, R.I., 1969.
11. A. W. Goodman, *Univalent Functions*, vols. I, II, Mariner, Tampa, FL, 1983.
12. K. Löwner, Untersuchungen über schlichte konforme Abbildungen des Einheitskreises, I, *Math. Ann.*, 89 (1923) 103–121.
13. I. M. Milin, *Univalent functions and orthonormal systems*. English transl., *Amer. Math. Soc.*, Providence, RI, 1977.
14. C. Pommerenke, *Univalent Functions*, Vandenhoeck and Ruprecht, Göttingen, 1975.
15. ———, The Bieberbach conjecture, *Math. Intelligencer*, 7 (1985) 23–25, 32.

NON-SEXIST SOLUTION OF THE MÉNAGE PROBLEM

KENNETH P. BOGART

Department of Mathematics, Dartmouth College, Hanover, NH 03755

PETER G. DOYLE

AT & T Bell Labs, Murray Hill, NJ 07974

1. The ménage problem. The *ménage problem* (problème des ménages) asks for the number M_n of ways of seating n man-woman couples at a circular table, with men and women

Kenneth Bogart received his B.S. in Mathematics from Marietta College in 1965 and his Ph.D. in Mathematics from CalTech in 1968. He has been teaching at Dartmouth College, where he is a professor of Mathematics and Computer Science, ever since. His research interests are in the applications of algebra to combinatorics, and the applications of combinatorics, especially to Computer Science. Much of his research has involved the theory or applications of ordered sets. His textbook, *Introductory Combinatorics*, was published in 1983 by Pitman Publishing Company. He is currently writing a textbook on discrete mathematics.

Peter Doyle got his Ph.D. in mathematics from Dartmouth College in 1982. While he was at Dartmouth, he and Laurie Snell wrote a book called *Random Walks and Electric Networks* which was published by the MAA in 1984 in the Carus Mathematical Monographs series. After getting his degree he worked as a systems programmer for a year, and spent a year at the Institute for Mathematics and its Applications in Minneapolis. He now works in the Mathematical Sciences Research Center at AT & T Bell Labs in Murray Hill, NJ.

alternating, so that no one sits next to his or her partner. This famous problem was initially posed by Lucas [8] in 1891, though an equivalent problem had been raised earlier by Tait [12] in connection with his work on knot theory (see Kaplansky and Riordan [6]). This problem has been discussed by numerous authors (see the references listed in [6]), and many solutions have been found. Most of these solutions tell how to compute M_n using recurrence relations or generating functions, as opposed to giving an explicit formula. The first explicit formula for M_n was published by Touchard [13] in 1934, though he did not give a proof. Finally, in 1943, Kaplansky [5] gave a proof of Touchard's formula. Kaplansky's derivation was simple but not quite straightforward, and the problem is still generally regarded to be tricky.

We will present a completely straightforward derivation of Touchard's formula. Like Kaplansky's, our solution is based on the principle of inclusion and exclusion (see Ryser [11] and Riordan [9]). What distinguishes our approach is that we do not seat the ladies (or gentlemen) first.

2. Solution to the relaxed ménage problem. We begin with an apparently simpler problem, called the *relaxed ménage problem*, which asks for the number m_n of ways of seating n couples around a circular table so that no one sits next to his or her partner. This is nearly the same as the ménage problem, only now we have relaxed the requirement that men and women alternate.

To determine m_n , we begin with the set S of all $(2n)!$ ways of seating the $2n$ individuals around the table, and use inclusion-exclusion on the set of couples who end up sitting together. Let us call the elements of S seatings, and let us denote by w_k the number of seatings under which some specified set of k couples (and possibly some other couples) end up sitting together. Clearly, w_k does not depend on the particular set of k couples we choose, and so, by the principle of inclusion and exclusion, we have

$$m_n = \sum_{k=0}^n (-1)^k \cdot \binom{n}{k} \cdot w_k.$$

To finish the enumeration, we must compute w_k . Assume $n > 1$. Let d_k denote the number of ways of placing k non-overlapping unlabeled dominos on $2n$ vertices arranged in a circle. (See Fig. 1.) Then

$$w_k = d_k \cdot k! \cdot 2^k \cdot (2n - 2k)!.$$




FIG. 1. Non-overlapping dominos.

(Decide where the k couples go, and which couple goes where, and which partner takes which seat, and where the $2n - 2k$ individuals go.) So now we have only to compute the d_k 's. This is a routine combinatorial problem. The answer is

$$d_k = \frac{2n}{2n-k} \cdot \binom{2n-k}{k}$$

(see Ryser [11], pp. 33–34, or Exercise 1 below). This yields

$$w_k = 2n \cdot (2n - k - 1)! \cdot 2^k.$$

Plugging this expression for w_k into the formula for m_n above, we get

$$m_n = \sum_{k=0}^n (-1)^k \cdot \binom{n}{k} \cdot 2n \cdot (2n - k - 1)! \cdot 2^k.$$

By symmetry, we know that m_n must be divisible by $2^n \cdot n!$. Pulling this factor out in front, we can write

$$m_n = 2^n \cdot n! \cdot \sum_{k=0}^n (-1)^k \cdot \frac{2n}{2n-k} \cdot \binom{2n-k}{k} \cdot (1 \cdot 3 \cdot 5 \cdots (2n - 2k - 1)).$$

The first few values of m_n are shown in Table 1.

TABLE 1. Relaxed ménage numbers.

n	m_n	$m_n/(2^n n!)$	$m_n/(2n)!$
2	8	1	0.333333...
3	192	4	0.266666...
4	11904	31	0.295238...
5	1125120	293	0.310052...
6	153262080	3326	0.319961...
7	28507207680	44189	0.326998...
8	6951513784320	673471	0.332246...
9	2153151603671040	11588884	0.336305...
10	826060810479206400	222304897	0.339537...

3. Solution to the ménage problem. For the ménage problem, we proceed just as before, only now we restrict the set S of seatings to those where men and women alternate. The number of these seatings is $2(n!)^2$: two ways to choose which seats are for men and which for women; $n!$ ways to seat the men in the men's seats; $n!$ ways to seat the women in the women's seats. Just as before, we have

$$M_n = \sum_{k=0}^n (-1)^k \cdot \binom{n}{k} \cdot W_k,$$

where W_k denotes the number of alternating seatings under which a specified set of k couples all end up sitting together. This time we have

$$W_k = 2 \cdot d_k \cdot k! \cdot (n - k)!^2.$$

(Decide which are men's seats and which women's, where the k couples go, which couple goes where, and where the $n - k$ men and $n - k$ women go.) Plugging in for d_k yields

$$W_k = 2 \cdot 2n \cdot (2n - k - 1)! \cdot \frac{(n - k)!^2}{(2n - 2k)!}.$$

Plugging this expression for W_k into the formula for M_n above, we get

$$M_n = \sum_{k=0}^n (-1)^k \cdot \binom{n}{k} \cdot 2 \cdot 2n \cdot (2n - k - 1)! \cdot \frac{(n - k)!^2}{(2n - 2k)!}.$$

By symmetry, we know that M_n must be divisible by $2 \cdot n!$. Pulling this factor out in front, we can write

$$M_n = 2 \cdot n! \cdot \sum_{k=0}^n (-1)^k \cdot \frac{2n}{2n-k} \cdot \binom{2n-k}{k} \cdot (n-k)!$$

The first few values of M_n are shown in Table 2.

TABLE 2. Ménage numbers.

n	M_n	$M_n/(2n!)$	$M_n/(2n!^2)$
2	0	0	0.0
3	12	1	0.166666...
4	96	2	0.083333...
5	3120	13	0.108333...
6	115200	80	0.111111...
7	5836320	579	0.114880...
8	382072320	4738	0.117509...
9	31488549120	43387	0.119562...
10	3191834419200	439792	0.121194...

4. Comparison with Kaplansky’s solution. The solution that we have just given is completely straightforward and elementary, yet we have said that the ménage problem is still generally regarded to be tricky. How can this be? The answer can be given in two words: “Ladies first.” It apparently never occurred to anyone who looked at the problem not to seat the ladies first (or in a few cases, the gentlemen). Thus Kaplansky and Riordan [6]: “We may begin by fixing the position of husbands and wives, say wives for courtesy’s sake.”

Seating the ladies first “reduces” the ménage problem to a problem of permutations with restricted position. Unfortunately, this new problem is more difficult than the problem we began with, as we may judge from the cleverness of Kaplansky’s solution [5]:

We now restate the *problème des ménages* in the usual fashion by observing that the answer is $2n!u_n$, where u_n is the number of permutations of $1, \dots, n$ which do not satisfy any of the following $2n$ conditions: 1 is 1st or 2nd, 2 is 2nd or 3rd, ..., n is n th or 1st. Now let us select a subset of k conditions from the above $2n$ and inquire how many permutations of $1, \dots, n$ there are which satisfy all k ; the answer is $(n - k)!$ or 0 according as the k conditions are compatible or not. If we further denote by v_k the number of ways of selecting k compatible conditions from the $2n$, we have, by the familiar argument of inclusion and exclusion, $u_n = \sum (-1)^k v_k (n - k)!$. It remains to evaluate v_k , for which purpose we note that the $2n$ conditions, when arrayed in a circle, have the property that only consecutive ones are not compatible...

Of course $v_k = d_k$, so we see how, by choosing to view the constraints as arrayed in a circle, Kaplansky has gotten back on the track of the straightforward solution. We can only admire Kaplansky’s cleverness in rediscovering the circle, and regret the tradition of seating the ladies first that made such cleverness necessary.

5. Conclusion. It appears that it was only the tradition of seating the ladies first that made the ménage problem seem in any way difficult. We may speculate that, were it not for this tradition, it would not have taken half a century to discover Touchard’s formula for M_n . Of all the ways in which sexism has held back the advance of mathematics, this may well be the most peculiar. (But see Exercise 2.)

6. Exercises. We list here, in the guise of exercises, some questions that you may want to explore with the help of the references listed.

1. Show how to “derive” the formula for d_k simply by writing down the answer, without using recurrence relations or generating functions or what have you. (Hint: Try this first for the formula for w_k .)
2. Was it really sexism that made the ménage problem appear difficult? (See Kaplansky and

Riordan [6], and the references listed there.)

3. Solve the analog of the *ménage* problem for the situation depicted in Figure 2. (No one is allowed to sit next to or across from his or her partner.)




FIG. 2. Real-world *ménage* problem.

4. Formulate the analog of the *ménage* problem for an arbitrary graph G , and show that it leads to a domino problem on G . Show that by seating the ladies or gentlemen first, and following Kaplansky’s lead, we arrive at a problem of how to place rooks on a chessboard. (See Riordan [9], Chap. 7.) Show that the domino problem and the rook problem are equivalent. Look into the relationship of the domino problem to the Ising model of statistical mechanics. (See Fisher [3], Kasteleyn [7].)

5. What problem was Tait [12] really interested in? Did Gilbert [4] solve it? Show that Gilbert could have used a simple Möbius inversion argument instead of Pólya’s theorem. What kinds of problems require the full force of Pólya’s theorem?

6. What does it mean to “solve” a combinatorial problem like the *ménage* problem? Is a closed-form solution better than a recurrence? What if what we really want is to generate configurations, rather than just count them? (See Wilf [14].)

7. Why did Tait not pursue the *ménage* problem? What do knots have to do with atomic spectra? What was it like to live in Nebraska in the 1880’s? (See Conway [2].)

8. The relaxed *ménage* problem can be further generalized as follows: Given two graphs G_1 and G_2 with the same number of vertices, find the number of one-to-one mappings of the vertices of G_1 onto the vertices of G_2 such that no pair of vertices that are adjacent in G_1 get sent to vertices that are adjacent in G_2 . Show that the dinner table problem (see Aspvall and Liang [1], Robbins [10]) can be phrased in these terms, and give a solution using inclusion-exclusion. Formulate and solve an “unrelaxed” version of this problem. Show that the *ménage* problem can be phrased in these terms, and discuss how useful this reformulation is. Do the same for the problem of enumerating Latin rectangles (see Ryser [11]).

References

1. B. Aspvall and F. M. Liang, The dinner table problem, Report STAN-CS-80-8222, Computer Science Department, Stanford University, Stanford, CA, 1980.
2. J. H. Conway, An enumeration of knots and links, and some of their algebraic properties, pp. 329–358 in *Computational Problems in Abstract Algebra*, ed. J. Leech, Pergamon, Oxford, 1970.
3. M. E. Fisher, Statistical mechanics of dimers on a plane lattice, *Phys. Rev.*, 124 (1961) 1664–1672.
4. E. N. Gilbert, Knots and classes of *ménage* permutations, *Scripta Math.*, 22 (1956) 228–233.
5. I. Kaplansky, Solution of the problème des ménages, *Bull. Amer. Math. Soc.*, 49 (1943) 784–785.
6. I. Kaplansky and J. Riordan, The problème des ménages, *Scripta Math.*, 12 (1946) 113–124.
7. P. W. Kasteleyn, Dimer Statistics and Phase Transitions, *J. Math. Phys.*, 4 (1963) 287–293.
8. E. Lucas, *Théorie des nombres*, Gauthier-Villars, Paris, 1891.
9. J. Riordan, *An Introduction to Combinatorial Analysis*, Wiley, New York, 1958.
10. D. Robbins, The probability that neighbors remain neighbors after random rearrangements, this *MONTHLY*, 87 (1980) 122–124.
11. H. J. Ryser, *Combinatorial Mathematics*, Carus Mathematical Monograph No. 14, Mathematical Association of America, Washington, DC, 1963.
12. P. G. Tait, On knots, I, II, III, pp. 273–347 in *Scientific Papers*, Cambridge Univ. Press, Cambridge, 1898.
13. J. Touchard, Sur un problème des permutations, *C. R. Acad. Sciences Paris*, 198 (1934) 631–633
14. H. Wilf, What is an answer?, this *MONTHLY*, 89 (1982) 289–292.