

- HW
- OFFICE HOURS
- RESERVE

Lecture

Group $\Sigma_3 = \langle e, (12), (123), (23), (13), (132) \rangle$

$G_1 = \left\langle \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \right\rangle G_1 \cong \Sigma_3 \leq GL_3(\mathbb{R})$

$G_2 = \left\langle \begin{pmatrix} -1 & -1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right\rangle G_2 \cong \Sigma_3$

two different representations of Σ_3

- Rep theory studies how to "represent" a group as matrices.
- Groups acting on vector spaces

Variations

Group: finite, discrete, algebraic
 Field: char 0 vs char p, alg closed?

More general: represent fin. dim. algs

Plan - Algebras For a While (I-IV)

- Sym groups, reps of sym g/cn.
- Current areas of research

Lecture 1

Review

Def R a ring (w/identity). A (left) R -module is an abelian group M together with an action $R \times M \rightarrow M, (r, m) \rightarrow rm$ such that

- $(r_1 + r_2)m = r_1m + r_2m$
- $r(m_1 + m_2) = rm_1 + rm_2$
- $(r_1 r_2)m = r_1(r_2m)$
- $1m = m$

$$\forall r, r_i \in R, m, m_i \in M$$

similarly for right R -modules.

Examples

1. Left regular module ${}_R R$ ($M = R$)
2. Given R -modules $\{M_i \mid i \in I\}$ then $\bigoplus_{i \in I} M_i$ is an R -module.
3. Any left ideal $I \subset R$, then I is a submodule.
4. Cat of \mathbb{Z} -modules equiv cat of abelian groups.

Def A submodule is a subgroup $N \subseteq M$ such that $rn \in N \forall r \in R, n \in N$.

A module is simple (aka irreducible) if the only submodules are zero and the entire module. INDECOMPOSABLE

Given a submodule $N \subseteq M$, the quotient group M/N is an R -module via $r(m+N) = rm+N$ ($r\bar{m} = \overline{rm}$), just check well-defined!

Thus no such thing as a "normal submodule".

* MODULE HOMOMORPHISMS, ISOMORPHISMS

Def. A composition series for an R -module M is a series of submodules

$0 = M_0 \subset M_1 \subset M_2 \dots \subset M_n = M$ such that M_i/M_{i-1} is simple.

A module satisfies the ACC if every ascending chain of submodules stops, and is called Noetherian. Similar D.C.C.

Jordan-Hölder Theorem Given any two series

$$0 = M_0 \subseteq \dots \subseteq M_r = M$$

$$0 = M'_0 \subseteq \dots \subseteq M'_s = M \quad \text{we may refine them to}$$

$$0 = L_0 \subseteq \dots \subseteq L_n = M$$

$0 = L'_0 \subseteq \dots \subseteq L'_n = M$ so that $\{L_i/L_{i-1}\}$ and $\{L'_i/L'_{i-1}\}$ are permutations, up to \cong . Thus TFAE

1. M has a composition series
2. Every series can be refined to a composition series
3. M satisfies ACC and DCC

Def. The length of a composition series is the composition length.
 M is uniserial if it has a unique composition series.

Ex $M = S_1 \oplus S_2$ is not!

Algebras

Remarks Our rings usually have additional vector space structure

Def R a field. A R -algebra is a vector space V over R which is also a ring, and operations are compatible, e.g. $\lambda(ab) = (\lambda a)b = a(\lambda b) \forall \lambda \in R, a, b \in V$.

- Remarks
1. Think of as a vector space w/ multiplication or as a ring w/ scalars
 2. Called a finite-dimensional algebra if f.d. as a vector space
 3. Usually assume identity, so $\{\lambda \cdot 1\}$ is a copy of the field inside.

Examples

1. Field R is a 1-dim R -algebra
2. Polynomial algebras $R[x_1, x_2, \dots, x_n]$ $>$ commutative
3. Matrix algebras $M_n(R) = \{n \times n \text{ matrices, entries in } R\}$
4. $T_n(R) = \left\{ \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \in M_n(R) \right\}$ subalgebra. (weaker than ideal)

5. Group algebras

G a group, RG is a vector space w/ G as a basis

Elements are finite linear combinations of group elements

• ~~dim~~ Multiplication is via distributive law

- RG has a distinguished basis, namely G
- Other elements are not necessarily invertible

• Occasionally "group rings" like $\mathbb{Z}G$, RG are useful.