

## Lecture 22

### Review

Let  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_t) \vdash d$ ,  $\lambda_t \neq 0$ . For any  $r \geq t$  we can represent  $\lambda$  on an abacus with  $r$  beads.

Then Let  $\lambda \vdash d$ . Then  $\lambda$  has a well-defined  $p$ -core  $\tilde{\lambda} \vdash d - pw$ , where  $w$  is the  $p$ -weight of  $\lambda$ . For  $\lambda, \mu \vdash d$ ,  $S^\lambda$  and  $S^\mu$  are in the same block iff  $\tilde{\lambda} = \tilde{\mu}$ .

### Recall

1.  $\lambda$  is uniquely determined by its  $p$ -core and  $p$ -quotient, a sequence of partitions  $\lambda_{(0)}, \lambda_{(1)}, \dots, \lambda_{(r)}$  of total weight  $w$ .
2. The # of irred  $\mathbb{C}\Sigma_d$  modules in a block is a function only of  $w$ . Similarly for  $k\Sigma_d$ .
3. Often use  $wp + t$  beads to represent  $\lambda$ .
4. The Defect group of a block of weight  $w$  is  $\cong$  to a Sylow  $p$ -subgroup of  $\Sigma_{pw}$ .
5. Adding/Removing nodes & rim  $p$ -hooks easy to represent on abacus.

Def Two categories  $\mathcal{C}$  &  $\mathcal{D}$  are equivalent if  $\exists$  functors  $\mathcal{F}: \mathcal{C} \rightarrow \mathcal{D}$  and  $\mathcal{G}: \mathcal{D} \rightarrow \mathcal{C}$  such that  $\mathcal{F} \circ \mathcal{G}$  is naturally  $\cong$  to  $\text{id}_{\mathcal{D}}$  and similarly  $\mathcal{G} \circ \mathcal{F} \cong \text{id}_{\mathcal{C}}$ .

Equivalently:  $\mathcal{F}: \mathcal{C} \rightarrow \mathcal{D}$  is an equivalence if for any objects  $G, G' \in \mathcal{C}$  the map  $\text{Hom}_{\mathcal{C}}(G, G') \rightarrow \text{Hom}_{\mathcal{D}}(\mathcal{F}(G), \mathcal{F}(G'))$  is a bijection and every object in  $\mathcal{D}$  is  $\cong$  to an object of form  $\mathcal{F}(G)$ .

Def Algebras  $A_1, A_2$  are Morita Equivalent if  $\text{mod } A_1$  and  $\text{mod } A_2$  are equivalent.

Ex  $R$  and  $M_n(R)$   $R$  a ring w  $1$ .

Donovan Conjecture Fix a  $p$ -group  $D$ .  $\exists$  only finitely many block algebras, up to Morita equivalence, with defect group  $\cong$  to  $D$ .

Thm (Scopes '91) Fix  $w \geq 0$ . There are only finitely many  $k\Sigma_d$  blocks of weight  $w$  up to Morita equivalence.

Moreover...


- # of non equiv blocks is  $\leq \prod_{i=1}^p ((i-1)(w-1) + 1)$

- Every Morita type occurs for some  $\Sigma_d$ ,

$$d \leq \frac{p^2(p-1)^2(w-1)^2}{4+wp}$$

### Cartan Matrix

Recall: Decomposition matrix  $D$  has



Def. Cartan matrix  $C$  has rows and columns indexed by irreducible  $RG$ -modules (equiv indec proj modules) and  $C_{ij}$

$$C_{ij} = \text{MULT}(P_i, D_j) = \text{Hom}(P_i, P_j) = \text{Hom}(P_i, P_j) = C_{ji}$$

Ex  $\Sigma_3$   $p=3$

$$\begin{matrix} S^3 \\ S^{21} \\ S^{13} \end{matrix} \begin{matrix} P^{20} & P^{21} \\ \begin{pmatrix} 1 & \\ & 1 \end{pmatrix} \\ \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \end{matrix}$$

PIMs

$R$	$S^n$
$S^n$	$R$
$R_1$	$S^n$

$$C = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$$

Thm  $C = D^t D$

Rmk Just as decomposition matrices of a block makes sense, so do Cartan matrices of a block.


Rmk Cartan matrix is invariant under Morita Equivalence (up to ordering)

### Scopes Idea

Fix a block  $B$  with  $p$ -core  $(b_1, b_{i-1}, b_i)$  and weight  $w$ .  
 Take an  $r+pw$  elt  $B$  set and repr. core on abacus


Let  $\Gamma_1, \Gamma_2, \dots, \Gamma_p$  be # of beads on each runner.

Ex  $p=3$ , core =  $(9, 7, 5, 3, 1^2)$



weight = 2  
 $r+pw = 12$

h.l. 1, 2, 5, 8, 11, 14



$\Gamma_1 = 2 \quad \Gamma_2 = 3 \quad \Gamma_3 = 7$

### Thm (Scopes)

Suppose for some  $i \geq 2$ ,  $\Gamma_i = \Gamma_{i-1} + k$  with  $k \geq w$ .  
 Let  $\bar{B}$  be block of  $\Sigma_{n-k}$  with core  $\bar{B}_i$  where  $\bar{B}$  has abacus obtained by swapping runners  $i-1$  and  $i$ .

Then  $B$  and  $\bar{B}$  are Morita equivalent.

Ex B block of  $\Sigma_{32}$  w/ core  $(9, 7, 5, 3, 1^2)$

$\bar{B}$

0 0 0  
 0 0 0  
 : 0 0  
 : 0 :  
 : 0 :  
 : 0 :  
 : 0 :  
 : 0 :

x x x x x x x  
 x x x x x x  
 x x x x  
 x x  
 x  
 x

block of  $\Sigma_{28}$  w/ core  $(8, 6, 4, 2)^2$

Repeat!

0 0 0  
 0 0 0  
 0 : 0  
 0 : :  
 0 : :  
 0 : :  
 0 : :  
 0 : :

x x x x x x  
 x x x x x  
 x x x  
 x  
 x

block of  $\Sigma_3$  core  $(7, 5, 3)^2$

Proof of Thm Assume Morita equivalences as above.  
 Suppose  $B, B'$  blocks of weight  $w$  of  $\Sigma_N, \Sigma_M$  w/  $N > M$ .  
 Suppose  $\exists B_0 = B', B_1, \dots, B_L = B$  so each pair Morita equiv as above.  
 Each family has a unique block ancestor of all in block.

Take such a block. Write out its 1<sup>st</sup> column hook lengths.  
 Since it's a  $p$ -core, 1<sup>st</sup> runner is empty.

EX. h.p. 1, 2, 5, 8, 11

0 0  
 : 0  
 : 0  
 : 0  
 : 0

Keep going

Let  $\theta_i$  be # of beads on each runner.

$\theta_1 = 0, \theta_i \leq \theta_{i-1} + w - 1$

Moreover Largest 1<sup>st</sup> col hook length is at most  $p(p-1)(w-1)$