

EXPANSIVE ACTIONS WITH SPECIFICATION OF SOFIC GROUPS, STRONG TOPOLOGICAL MARKOV PROPERTY, AND SURJUNCTIVITY

TULLIO CECCHERINI-SILBERSTEIN, MICHEL COORNAERT, AND HANFENG LI

To Klaus Schmidt on his 80th anniversary

ABSTRACT. A dynamical system is a pair (X, G) , where X is a compact metrizable space and G is a countable group acting by homeomorphisms of X . An endomorphism of (X, G) is a continuous selfmap of X which commutes with the action of G . One says that a dynamical system (X, G) is surjunctive provided that every injective endomorphism of (X, G) is surjective (and therefore is a homeomorphism). We show that when G is sofic, every expansive dynamical system (X, G) with nonnegative sofic topological entropy and satisfying the weak specification and the strong topological Markov properties, is surjunctive.

CONTENTS

1. Introduction	2
2. Preliminaries	5
2.1. General notation	5
2.2. Actions	6
2.3. Shifts and subshifts	6
2.4. Dynamical systems	6
2.5. Expansivity	7
2.6. The strong topological Markov property	7
2.7. The weak specification property	8
2.8. The pseudo-orbit tracing property	9
2.9. Algebraic dynamical systems	9
2.10. The Hamming distance	10
2.11. Sofic groups	10
2.12. Sofic topological entropy	11
3. Proofs	13
4. Examples	19
Appendix A. Stirling's approximation formula	20

Date: February 9, 2024.

2010 Mathematics Subject Classification. 37B40, 37B10, 37D20, 20F65.

Key words and phrases. Sofic group, sofic entropy, surjunctive dynamical system, expansive action, strong topological Markov property, weak specification property, subshift, surjunctive subshift, strongly irreducible subshift, algebraic dynamical system.

1. INTRODUCTION

Let A be a finite set, called the *alphabet*, and let G be a group. We equip $A^G = \prod_{g \in G} A$ with the *prodiscrete* topology, i.e., the product topology obtained by taking the discrete topology on each factor A of A^G , and the *shift* action of G , given by $gx(h) := x(g^{-1}h)$ for all $g, h \in G$ and $x \in A^G$. This action is clearly continuous. A closed G -invariant subset X of A^G is called a *subshift* and a continuous map $f: X \rightarrow X$ which commutes with the shift is called a *cellular automaton* (cf. [10]).

A group G is called *surjunctive* if, given any finite alphabet A , every injective cellular automaton $f: A^G \rightarrow A^G$ is surjective (and therefore a homeomorphism) (see [10, Chapter 3]). Lawton [28] proved that all residually finite groups are surjunctive and Gottschalk [22] raised the question of determining those groups which are surjunctive. The statement asserting that every group is surjunctive is now commonly known as the *Gottschalk conjecture*. The fact that all amenable groups are surjunctive follows from one of the implications of the Garden of Eden theorem established in [14] (see also [10, Chapter 5]). Up to now, the largest class of groups that are known to be surjunctive is the class of *sofic* groups introduced by Gromov in [23] and named by Weiss in [36] (see Section 2.11): surjunctivity of sofic groups was established by Gromov [23] and Weiss [36] (see also [8], [10], [25] for expositions of this result). The class of sofic groups contains all residually finite groups, all amenable groups, and, more generally, all residually amenable groups. Actually, no example of a non-sofic group has yet been found.

A *dynamical system* is a pair (X, G) consisting of a compact metrizable space X equipped with a continuous action of a countable group G . An *endomorphism* of a dynamical system (X, G) is a continuous G -equivariant map $f: X \rightarrow X$.

For instance, if A is a finite set, G is a countable group, and $X \subset A^G$ is a subshift, then (X, G) , where X is equipped with the topology induced by the prodiscrete topology and the action of G obtained by restriction of the shift action, is a dynamical system whose endomorphisms are the cellular automata $f: X \rightarrow X$.

One says that a dynamical system (X, G) is *surjunctive* if every injective endomorphism of (X, G) is surjective (and hence a homeomorphism). Also (X, G) is *expansive* if, given a compatible metric ρ on X , there exists a constant $c > 0$ such that for all distinct $x, y \in X$ there exists $g \in G$ such that $\rho(gx, gy) > c$. By compactness of X , the fact that the dynamical system (X, G) is expansive or not does not depend on the choice of the compatible metric ρ . For instance, if $X \subset A^G$ is a subshift, then (X, G) is expansive. See Section 2.5.

A *sofic approximation* of a countable group G is a sequence $\Sigma = (d_j, \sigma_j)_{j \in \mathbb{N}}$, where d_j is a positive integer and $\sigma_j: G \rightarrow \text{Sym}(d_j)$ is a map from G into the symmetric group of degree d_j , satisfying the following conditions:

- (i) $\lim_{j \rightarrow \infty} d_j = +\infty$,
- (ii) $\lim_{j \rightarrow \infty} \eta_{d_j}(\sigma_j(s)\sigma_j(t), \sigma_j(st)) = 0$ for all $s, t \in G$, and

(iii) $\lim_{j \rightarrow \infty} \eta_{d_j}(\sigma_j(s), \sigma_j(t)) = 1$ for all distinct $s, t \in G$, where η_{d_j} denotes the normalized *Hamming distance* on $\text{Sym}(d_j)$. A countable group G is *sofic* if it admits a sofic approximation. More generally, a group is *sofic* if all of its finitely generated subgroups are sofic.

For sofic groups, David Kerr and the third named author [24, 25, 26] defined and studied the *topological entropy* $h_\Sigma(X, G)$ of a dynamical system (X, G) relative to a sofic approximation Σ of the group G . See Section 2.12.

The notion of *specification*, a strong orbit tracing property, was introduced by Rufus Bowen for \mathbb{Z} -actions in relation to his studies on Axiom A diffeomorphisms in [5] (see also [19, Definition 21.1]) and was subsequently extended to \mathbb{Z}^d -actions by Ruelle in [34]. Several versions and generalizations of specification have appeared in the literature (see, in particular, [30, Definition 5.1] and [16, Definition 6.1]). Here is the one we need (cf. [16, Definition 6.1], see also [29]). A dynamical system (X, G) has the *weak specification property* if for any $\varepsilon > 0$ there exists a finite symmetric subset $F \subset G$ containing 1_G satisfying the following property: if $(\Omega_i)_{i \in I}$ is any finite family of finite subsets of G such that $F\Omega_i \cap \Omega_j = \emptyset$ for all distinct $i, j \in I$, and $(x_i)_{i \in I}$ is any family of points in X , then there exists $x \in X$ such that $\rho(sx, sx_i) \leq \varepsilon$ for all $i \in I$ and $s \in \Omega_i$, where ρ is any metric compatible with the topology on X . When $X \subset A^G$ is a subshift, the dynamical system (X, G) has the weak specification property if and only if X is *strongly irreducible* (cf. [9], [29, Proposition A.1]). See Section 2.7.

In [2], Barbieri, García-Ramos, and the third named author, inspired by investigations in the context of symbolic dynamics [3], introduced and studied various notions of topological Markov properties. Among them, the following will play a crucial role in our work. A dynamical system (X, G) satisfies the *strong topological Markov property* (briefly, *sTMP*) if for any $\varepsilon > 0$ there exists $\delta > 0$ and a finite subset $F \subset G$ containing 1_G such that the following holds: given any finite subset $A \subset G$ and $x, y \in X$ such that $\rho(sx, sy) \leq \delta$ for all $s \in FA \setminus A$, there exists $z \in X$ satisfying that $\rho(sz, sx) \leq \varepsilon$ for all $s \in FA$ and $\rho(sz, sy) \leq \varepsilon$ for all $s \in G \setminus A$. When $X \subset A^G$ is a subshift, the strong topological Markov property for (X, G) is equivalent to X being *splicable* in the sense of Gromov [23, Section 8.C] (see also [13, Section 3.8]). For instance, every subshift of *finite type* is splicable. See Section 2.6.

We are now in a position to state the main result of the paper.

Theorem 1.1. *Let G be a countable sofic group and let (X, G) be an expansive dynamical system with the weak specification property and the strong topological Markov property. Assume that $h_\Sigma(X, G) \geq 0$ for some sofic approximation Σ of G . Then (X, G) is surjective.*

The key ingredient for the proof of Theorem 1.1 is the following simple classical argument: a dynamical system (X, G) , admitting a topological conjugacy invariant taking values in an ordered set which are strictly smaller than that of (X, G) on proper subsystems, is surjective. The following result, which is of independent interest, shows that (in the hypotheses of Theorem 1.1) sofic topological entropy yields such a suitable topological conjugacy invariant:

Proposition 1.2. *Let G be a countable sofic group and let (X, G) be an expansive dynamical system with the weak specification property and the strong topological Markov property. Assume that $h_\Sigma(X, G) \geq 0$ for some sofic approximation Σ of G . Suppose that $Y \subset X$ is a proper closed G -invariant subset. Then $h_\Sigma(Y, G) < h_\Sigma(X, G)$.*

As an immediate consequence of Theorem 1.1 we deduce the following:

Corollary 1.3. *Let A be a finite set and let G be a countable sofic group. Suppose that $X \subset A^G$ is a splicable (e.g., of finite type) strongly irreducible subshift. Assume that $h_\Sigma(X, G) \geq 0$ for some sofic approximation Σ of G . Then (X, G) is surjective. \square*

As the full shift A^G is (trivially) splicable (in fact of finite type) and strongly irreducible, and $h_\Sigma(A^G, G) = \log |A| \geq 0$ for all sofic approximations Σ of any sofic group G , from Corollary 1.3 one immediately recovers the Gromov-Weiss surjectivity theorem for sofic groups. When G is an amenable group, the surjectivity result in Corollary 1.3 was proved under the additional assumption of X being of finite type by Fiorenzi [21, Corollary 4.8]; the first two named authors [9] then showed that the strong irreducibility of the subshift is sufficient to imply surjectivity when G is amenable. Note that, however, the strong irreducibility condition cannot be dropped: for instance, Weiss [36] gave an example of a subshift of finite type over $G = \mathbb{Z}$ which is not surjective (see Example 4.2) as well as an example of a topologically mixing subshift of finite type over $G = \mathbb{Z}^2$ which is not surjective either.

An *algebraic dynamical system* is a dynamical system (X, G) , where X is a compact metrizable abelian group and G is a countable group acting by continuous automorphisms of X . Given an algebraic dynamical system (X, G) , the Pontryagin dual \widehat{X} , when equipped with the dual action of G , is a countable left $\mathbb{Z}[G]$ -module (see Section 2.9). In fact, Pontryagin duality establishes a bijective correspondence between algebraic dynamical systems over G and countable left $\mathbb{Z}[G]$ -modules.

An algebraic dynamical system (X, G) is said to be *finitely presented* provided its Pontryagin dual \widehat{X} is a finitely presented left $\mathbb{Z}[G]$ -module. This amounts to saying that there exists an $n \times m$ matrix A with coefficients in $\mathbb{Z}[G]$ such that $\widehat{X} = \mathbb{Z}[G]^m / (\mathbb{Z}[G]^n A)$; we then denote X by X_A . In the particular case when A is a square matrix, the associated algebraic dynamical system (X_A, G) is called a *generalized principal algebraic dynamical system*. Note that if (X, G) is an algebraic dynamical system, then G fixes the identity element of X . Therefore, if in addition the group G is sofic, then (X, G) satisfies $h_\Sigma(X, G) \geq 0$ for all sofic approximations Σ of G . From Theorem 1.1, in combination with results of Meyerovitch [31], Ren [33], and Barbieri, García-Ramos and the third named author [2], we then deduce the following:

Corollary 1.4. *Let G be a countable sofic group. Let $A \in \text{Mat}_n(\mathbb{Z}[G])$ and suppose that A is invertible in $\text{Mat}_n(\ell^1(G))$. Then the generalized principal algebraic dynamical system (X_A, G) is surjective.*

Here $\ell^1(G) := \{f \in \mathbb{R}^G : \sum_{g \in G} |f(g)| < \infty\}$ is the Banach algebra of all real summable functions on G . Note that when $n = 1$ and $m \geq 2$ is an integer, then taking $f = m1_G \in$

$\mathbb{Z}[G]$ one has $X_f = \{1, 2, \dots, m\}^G$, the full shift over an m -elements alphabet set. From Corollary 1.4 one then recovers, once more, the Gromov-Weiss surjunctivity theorem for sofic groups.

We don't know if the above corollary can be extended to all expansive finitely presented algebraic dynamical systems.

Surjunctivity of algebraic dynamical systems, in relation with the so-called topological rigidity property, was investigated also in [4].

We conclude with a surjunctivity result for actions of amenable groups on (not necessarily abelian) compact metrizable groups by continuous automorphisms. First, recall that given a countable amenable group G , a dynamical system (X, G) has *completely positive entropy* (briefly *CPE*) if every nontrivial factor has positive topological entropy. We then have the following result whose proof is not a consequence of Proposition 1.2 (or Theorem 1.1) but can be read independently.

Theorem 1.5. *Let G be a countably infinite amenable group G and let (X, G) be a dynamical system, where X is a compact metrizable group and G acts by continuous automorphisms of X . Suppose that (X, G) has CPE and that $h_{\text{top}}(X, G) < \infty$ (e.g., (X, G) is expansive). Then (X, G) is surjunctive.*

Acknowledgments. We express our gratitude to Felipe García-Ramos and to the anonymous referee for their several interesting remarks. H. Li wishes to thank Filippo Tolli (Università Roma Tre) and the Department SBAI of the Sapienza University in Rome for hospitality during a visit in December 2019, in which this work started. He was partially supported by NSF grant DMS-1900746.

2. PRELIMINARIES

In this section we fix the notation and review some background material on dynamical systems, sofic groups, and sofic entropy.

2.1. General notation. Throughout the paper, we use $|\cdot|$ to denote cardinality of finite sets.

We write \mathbb{N} for the set of non-negative integers.

Given sets A and B , we denote by A^B the set consisting of all maps $x: B \rightarrow A$. We will sometimes regard an element $x \in A^B$ as a family of elements of A indexed by B , namely $x = (x(b))_{b \in B}$, where $x_b := x(b)$ for all $b \in B$. If $B' \subset B$ and $x \in A^B$, we denote by $x|_{B'} \in A^{B'}$ the *restriction* of x to B' , that is, the map $x|_{B'}: B' \rightarrow A$ defined by $x|_{B'}(b') := x(b')$ for all $b' \in B'$.

Given a set X , we denote by $\text{Sym}(X)$ the *symmetric group* of X , i.e., the group consisting of all bijective maps $\sigma: X \rightarrow X$ with the composition of maps as the group operation.

Given a positive integer d , we write $[d] := \{1, 2, \dots, d\}$. To simplify notation, we write $\text{Sym}(d)$ instead of $\text{Sym}([d])$.

Given a group G , we denote by $\mathcal{F}(G)$ the set of all nonempty finite subsets $F \subset G$. Also, given $F \in \mathcal{F}(G)$, we say that a (not necessarily finite) subset $V \subset G$ is F -separated if $Fs \cap Ft = \emptyset$ for all distinct $s, t \in V$.

Given a metric space (X, ρ) and a subset $Y \subset X$, we denote by $\text{diam}(Y, \rho) := \sup\{\rho(y_1, y_2) : y_1, y_2 \in Y\}$ the *diameter* of Y . For $x \in X$, we write $\rho(x, Y) := \inf_{y \in Y} \rho(x, y)$. Moreover, given $\varepsilon > 0$ one says that Y is (ρ, ε) -separated provided that $\rho(y, y') \geq \varepsilon$ for all distinct $y, y' \in Y$.

2.2. Actions. An *action* of a group G on a set X is a group morphism $\alpha: G \rightarrow \text{Sym}(X)$. If α is an action of a group G on a set X , given $g \in G$ and $x \in X$, we usually write $\alpha_g(x)$ or, if the action is clear from the context, gx , instead of $\alpha(g)(x)$.

An action of a group G on a topological space X is said to be continuous if its image is contained in the group of homeomorphisms of X .

2.3. Shifts and subshifts. Let A be a finite set and let G be a countable group. We equip A with the discrete topology and $A^G = \prod_{g \in G} A$ with the product topology (this is also called the *prodiscrete topology* on A^G). Note that A^G is a compact space (by Tychonoff's theorem) and is metrizable since G is countable. A metric inducing the prodiscrete topology can be constructed as follows. Let $(\Omega_n)_{n \in \mathbb{N}}$ be a sequence of finite subsets of G satisfying that $\Omega_0 = \emptyset$, $\Omega_1 = \{1_G\}$, $\Omega_n^{-1} := \{g^{-1} : g \in \Omega_n\} = \Omega_n$, $\Omega_n \subset \Omega_{n+1}$ for all $n \in \mathbb{N}$, and $\bigcup_{n \in \mathbb{N}} \Omega_n = G$. Then the map $\rho: A^G \times A^G \rightarrow [0, 1]$ defined by setting

$$(2.1) \quad \rho(x, y) := 2^{-n(x, y)}, \text{ where } n(x, y) := \sup\{n \in \mathbb{N} : x|_{\Omega_n} = y|_{\Omega_n}\}$$

(with the usual convention that $2^{-\infty} = 0$) for all $x, y \in A^G$, is a metric on A^G inducing the prodiscrete topology.

We make G act on A^G by the *shift* action, given by the map $G \times A^G \rightarrow A^G$, $(g, x) \mapsto gx$, where

$$(2.2) \quad gx(h) := x(g^{-1}h) \text{ for all } g, h \in G \text{ and } x \in A^G.$$

The action of G on A^G is clearly continuous.

A closed G -invariant subset of A^G is called a *subshift*.

A subshift $X \subset A^G$ is said to be of *finite type* provided there exists a finite subset $\Omega \subset G$ and $\mathcal{A} \subset A^\Omega$ such that $X = \{x \in A^G : (gx)|_\Omega \in \mathcal{A} \text{ for all } g \in G\}$ (such a set \mathcal{A} is then called a *defining set of admissible patterns* for X and the subset Ω is called a *memory set* for X).

2.4. Dynamical systems. A *dynamical system* is a pair (X, G) consisting of a compact metrizable space X , called the *phase space*, equipped with a continuous action of a countable group G .

Let (X, G) be a dynamical system and let ρ be a metric on X compatible with the topology. Given a nonempty subset $A \subset G$, we denote by ρ_A the metric on X defined by setting

$$(2.3) \quad \rho_A(x, y) = \sup_{g \in A} \rho(gx, gy)$$

for all $x, y \in X$. Note that $\rho_{\{1_G\}} = \rho$.

2.5. Expansivity.

Definition 2.1. The dynamical system (X, G) is *expansive* if there exists a constant $c > 0$ such that for all distinct $x, y \in X$ there exists $g \in G$ such that $\rho(gx, gy) > c$. Such a value c is called an *expansivity constant* for (X, G) (relative to the metric ρ).

Compactness of the phase space X guarantees that the definition of expansivity does not depend on the choice of the metric ρ . Note that $c > 0$ is an expansivity constant for (X, G) if and only if $\rho_G(x, y) > c$ for all distinct $x, y \in X$ (cf. (2.3)). For example, if A is a finite set and G is a countable group, then the symbolic dynamical system (A^G, G) is expansive (with expansivity constant $c = 1/2$ relative to the metric ρ defined by (2.1)).

The following lemma (cf. [2, Lemma 3.13]) was proved by Bryant [7, Theorem 5] (cf. [1, Proposition 2]) for $G = \mathbb{Z}$.

Lemma 2.2 (Uniform expansivity). *Let (X, G) be a dynamical system, let ρ be a compatible metric on X , and let $c > 0$. The following conditions are equivalent:*

- (a) (X, G) is expansive and c is an expansivity constant for (X, G) relative to ρ ;
- (b) for every $\varepsilon > 0$, there exists a nonempty finite subset $K \subset G$ such that the following holds: if $x, y \in X$ satisfy that $\rho_K(x, y) \leq c$, then $\rho(x, y) < \varepsilon$.

Proof. Assume (a) and suppose, by contradiction, that (b) fails to hold, that is, there exists $\varepsilon_0 > 0$ such that for every nonempty finite subset $K \subset G$ there exist $x_K, y_K \in X$ such that $\rho_K(x_K, y_K) \leq c$ but $\rho(x_K, y_K) \geq \varepsilon_0$. Consider the nets $(x_K)_{K \in \mathcal{F}(G)}$ and $(y_K)_{K \in \mathcal{F}(G)}$ of points in X , where the common index set $\mathcal{F}(G)$ is partially ordered by inclusion. Since X is compact, we can find a cluster point (x, y) in $X \times X$ for $(x_K, y_K)_{K \in \mathcal{F}(G)}$. Note that, by continuity, $\rho(x, y) \geq \varepsilon_0$, so that, in particular, $x \neq y$. Let $g \in G$. If $K \in \mathcal{F}(G)$ contains g one has $\rho(gx_K, gy_K) \leq c$. By continuity, we have $\rho(gx, gy) \leq c$. As g was arbitrary, we deduce that $\rho_G(x, y) \leq c$. Since $x \neq y$, this contradicts the assumption that c is an expansivity constant. This proves the implication (a) \Rightarrow (b).

Conversely, suppose (b). Let $x, y \in X$ such that $\rho_G(x, y) \leq c$ and let us show that $x = y$. Let $\varepsilon > 0$ and let $K \subset G$ be a finite subset as in (b). As $\rho_K(x, y) \leq \rho_G(x, y) \leq c$, we have $\rho(x, y) < \varepsilon$. Since ε was arbitrary, this implies that $\rho(x, y) = 0$, that is, $x = y$. This shows that c is an expansivity constant for (X, G) . \square

2.6. The strong topological Markov property.

Definition 2.3. The dynamical system (X, G) satisfies the *strong topological Markov property* (briefly, *sTMP*) if for any $\varepsilon > 0$ there exists $\delta > 0$ and a finite subset $F \subset G$ containing 1_G such that the following holds. Given any finite subset $A \subset G$ and $x, y \in X$ such that $\rho_{FA \setminus A}(x, y) \leq \delta$, there exists $z \in X$ satisfying that $\rho_{FA}(z, x) \leq \varepsilon$ and $\rho_{G \setminus A}(z, y) \leq \varepsilon$.

Definition 2.4. The dynamical system (X, G) satisfies the *uniform strong topological Markov property* if for any $\varepsilon > 0$ there exists $\delta > 0$ and a finite subset $F \subset G$ containing 1_G such that the following holds. Given a finite subset $A \subset G$, an FA -separated subset

$V \subset G$, a family $(x_v)_{v \in V}$ of points in X , and $y \in X$ such that $\rho_{(FA \setminus A)s}(x_s, y) \leq \delta$ for all $s \in V$, there exists $z \in X$ satisfying that $\rho_{FA_s}(x_s, z) \leq \varepsilon$ for all $s \in V$ and $\rho_{G \setminus AV}(y, z) \leq \varepsilon$.

Remark 2.5. Clearly, every dynamical system with uniform sTMP satisfies sTMP (just take $V = \{1_G\}$). Conversely, it is shown in [2, Remark 3.14] that every expansive dynamical system satisfying sTMP satisfies the uniform sTMP.

In the symbolic setting, we have the following characterization of sTMP and usTMP (see [2, Corollary 3.8] and [13, Proposition 3.16]).

Proposition 2.6. *Let A be a finite set, let G be a countable group, and let $X \subset A^G$ be a subshift. Then the following conditions are equivalent:*

- (a) (X, G) satisfies the strong topological Markov property;
- (b) (X, G) satisfies the uniform strong topological Markov property;
- (c) *there exists a finite subset $S \subset G$ containing 1_G such that the following holds. Given any finite subset $A \subset G$ and $x, y \in X$ such that $x|_{AS \setminus A} = y|_{AS \setminus A}$, the configuration $z \in A^G$ such that $z|_A = x|_A$ and $z|_{G \setminus A} = y|_{G \setminus A}$ belongs to X .*

Moreover, a sufficient condition for the above equivalent conditions is X being of finite type. \square

Remark 2.7. A subshift $X \subset A^G$ satisfying condition (c) (equivalently, conditions (a) and (b)) in Proposition 2.6 was termed *splicable* by Gromov in [23, Section 8.C] (see also [13, Section 3.8]).

2.7. The weak specification property.

Definition 2.8. The dynamical system (X, G) satisfies the *weak specification property* (briefly, *wSP*) if for any $\varepsilon > 0$ there exist a finite symmetric subset $F \subset G$ containing 1_G such that the following holds. Given finite subsets $A_1, A_2, \dots, A_n \subset G$ satisfying $FA_i \cap FA_j = \emptyset$ for all distinct $i, j = 1, 2, \dots, n$, and $x_1, x_2, \dots, x_n \in X$, there exists $z \in X$ such that $\rho_{A_i}(x_i, z) \leq \varepsilon$ for all $i = 1, 2, \dots, n$.

Remark 2.9. If wSP holds, then one may allow the subsets $A_1, A_2, \dots, A_n \subset G$ in Definition 2.8 to be infinite.

In the symbolic setting, the wSP has the following characterization (see [29, Proposition A.1] and [11, Proposition 6.7]).

Proposition 2.10. *Let A be a finite set, let G be a countable group, and let $X \subset A^G$ be a subshift. Then the following conditions are equivalent:*

- (a) *the dynamical system (X, G) satisfies the weak specification property;*
- (b) *there exists a finite subset $S \subset G$ containing 1_G such that the following holds. Given finite subsets $A_1, A_2 \subset G$ such that $A_1 S \cap A_2 = \emptyset$ and $x_1, x_2 \in X$, there exists $z \in X$ such that $z|_{A_1} = x_1|_{A_1}$ and $z|_{A_2} = x_2|_{A_2}$.*

\square

Remark 2.11. A subshift satisfying condition (b) (equivalently, condition (a)) in Proposition 2.10 is called *strongly irreducible* (cf. [21], [9]).

We mention that M. Doucha [20] recently showed that any expansive action of a countable amenable group G on a compact metrizable space X satisfying both the weak specification and the strong topological Markov properties also satisfies the so-called *Moore property*, i.e. every surjective endomorphism of (X, G) is *pre-injective* (the latter means that the restriction to each homoclinicity class of (X, G) is injective). This, together with an earlier result of the third-named author [29] (where the strong topological Markov property is not needed), namely, the so-called *Myhill property*, i.e. every pre-injective endomorphism of (X, G) is surjective, establishes the Garden of Eden theorem for all expansive actions of countable amenable groups on compact metrizable spaces satisfying the weak specification and the strong topological Markov properties.

2.8. The pseudo-orbit tracing property. The pseudo-orbit tracing property (briefly, POTP) was originally introduced and studied by R. Bowen [6] for \mathbb{Z} -actions, motivated by the study of Axiom A diffeomorphisms. Walters [35] studied the relationships between POTP and topological stability and proved, among other things, that every topologically stable homeomorphism of a compact manifold of dimension ≥ 2 has the POTP. Chung and Lee [15] recently considered the pseudo-orbit tracing property for actions of (finitely generated) countable groups and extended Walters's results. In particular, they proved that every expansive dynamical system (X, G) with the POTP is topologically stable. Here is the definition:

Definition 2.12. Let (X, G) be a dynamical system and let ρ be a compatible metric on X . Given $\delta > 0$ and a finite subset $S \subset G$ an (S, δ) *pseudo-orbit* is a sequence $(x_g)_{g \in G}$ in X such that $\rho(sx_g, x_{sg}) < \delta$ for all $s \in S$ and $g \in G$. Then one says that (X, G) has the *pseudo-orbit tracing property* if for every $\varepsilon > 0$ there exists $\delta > 0$ and a finite subset $S \subset G$ such that for every (S, δ) pseudo-orbit $(x_g)_{g \in G}$ in X there exists $x \in X$ such that $\rho(gx, x_g) < \varepsilon$ for all $g \in G$ (one then says that the pseudo-orbit is ε -traced by x).

Barbieri, García-Ramos, and the third-named author [2, Proposition 3.10] proved that every dynamical system with the pseudo-orbit tracing property has the strong topological Markov property.

2.9. Algebraic dynamical systems. Let L be a locally compact Abelian group. The *Pontryagin dual* of L is the set \widehat{L} of all continuous group homomorphism $\chi: L \rightarrow \mathbb{T} = \mathbb{R}/\mathbb{Z}$, called the *characters* of L . When equipped with the topology given by uniform convergence on compact sets, \widehat{L} is a locally compact Abelian group as well. Moreover, if L is compact (resp. discrete, resp. countable and discrete) then \widehat{L} is discrete (resp. compact, resp. compact metrizable). The so-called *Pontryagin duality* establishes that the *evaluation map* $\text{ev}_L: L \rightarrow \widehat{\widehat{L}}$, defined by setting $\text{ev}_L(\ell)(\chi) := \chi(\ell)$ for all $\ell \in L$ and $\chi \in \widehat{L}$, yields a canonical isomorphism (of locally compact topological groups) of L into its bi-dual $\widehat{\widehat{L}}$. (We refer to [32] for more on Pontryagin duality.)

Let G be a countable group and let $\mathbb{Z}[G]$ denote the integral group ring of G . Suppose that G acts on L by continuous group automorphisms. Then the action of G induces an action on the Pontryagin dual \widehat{L} by continuous group automorphisms and, by linearity,

a left $\mathbb{Z}[G]$ -module structure on both L and \widehat{L} . In particular, the evaluation map is G -equivariant.

An *algebraic dynamical system* is a pair (X, G) , where X is a compact metrizable Abelian topological group and G is a countable group acting on X by continuous group automorphisms.

Let M be a countable left $\mathbb{Z}[G]$ -module. Then, if we equip the Abelian group M with its discrete topology, its Pontryagin dual \widehat{M} is a compact metrizable Abelian group and (\widehat{M}, G) is an algebraic dynamical system. In this way, algebraic dynamical systems with acting group G are in one-to-one correspondence with countable left $\mathbb{Z}[G]$ -modules.

An algebraic dynamical system (X, G) is said to be *finitely presented* provided its Pontryagin dual \widehat{X} is a finitely presented left $\mathbb{Z}[G]$ -module. For a ring R we denote by $\text{Mat}_{n,k}(R)$ the left R -module of all $n \times k$ matrices with coefficients in R . If $n = k$ we simply write $\text{Mat}_n(R)$ instead of $\text{Mat}_{n,k}(R)$. Let $k, n \in \mathbb{N}$ and let $A \in \text{Mat}_{n,k}(\mathbb{Z}[G])$. Then $\mathbb{Z}[G]^n A$ is a finitely generated left $\mathbb{Z}[G]$ -submodule of $\mathbb{Z}[G]^k$. We denote by $M_A := \mathbb{Z}[G]^k / (\mathbb{Z}[G]^n A)$ the corresponding finitely presented left $\mathbb{Z}[G]$ -module and by $X_A := \widehat{M_A}$ its Pontryagin dual. The algebraic dynamical system (X_A, G) is called the *finitely presented algebraic dynamical system* associated with A . An algebraic dynamical system (X, G) is finitely presented if and only if there exist $k, n \in \mathbb{N}$ and $A \in \text{Mat}_{n,k}(\mathbb{Z}[G])$ such that (X, G) is *isomorphic* to (X_A, G) (i.e., there exists a continuous G -equivariant group isomorphism $X \rightarrow X_A$).

An algebraic dynamical system (X, G) is said to be *principal* if \widehat{X} is a cyclic left $\mathbb{Z}[G]$ -module whose annihilator is a principal left ideal of $\mathbb{Z}[G]$. This amounts to saying that (X, G) is isomorphic to (X_f, G) for some $f \in \mathbb{Z}[G]$, where (X_f, G) is the algebraic dynamical system whose Pontryagin dual is $\mathbb{Z}[G]/\mathbb{Z}[G]f$ (here $k = n = 1$).

2.10. The Hamming distance. Let d be a positive integer. The map $\eta_d: \text{Sym}(d) \times \text{Sym}(d) \rightarrow [0, 1]$ defined by setting

$$(2.4) \quad \eta_d(\sigma, \sigma') := \frac{1}{d} |\{a \in [d] : \sigma(a) \neq \sigma'(a)\}|$$

for all $\sigma, \sigma' \in \text{Sym}(d)$, is a bi-invariant metric on $\text{Sym}(d)$, called the *Hamming distance*.

2.11. Sofic groups. Let G be a countable group. A *sofic approximation* of G is a sequence $\Sigma = (d_j, \sigma_j)_{j \in \mathbb{N}}$, such that the following conditions are satisfied:

- d_j is a positive integer and $\lim_{j \rightarrow \infty} d_j = +\infty$;
- $\sigma_j: G \rightarrow \text{Sym}(d_j)$ is a map from G into the symmetric group of $[d_j]$;
- $\lim_{j \rightarrow \infty} \eta_{d_j}(\sigma_j(st), \sigma_j(s)\sigma_j(t)) = 0$ for all $s, t \in G$;
- $\lim_{j \rightarrow \infty} \eta_{d_j}(\sigma_j(s), \sigma_j(t)) = 1$ for all distinct $s, t \in G$,

where η_{d_j} denotes the Hamming distance on $\text{Sym}(d_j)$ defined by Formula (2.4).

One says that a (possibly uncountable) group is *sofic* if each of its finitely generated subgroups admits a sofic approximation.

2.12. Sofic topological entropy. In this section we recall the definition of the sofic topological entropy introduced by Kerr and the third-named author in [26, Chapter 10].

Let G be a countable sofic group and let (X, G) be a dynamical system. We fix a metric ρ on X compatible with the topology and a sofic approximation $\Sigma = (d_j, \sigma_j)_{j \in \mathbb{N}}$ of G .

Let $d \geq 1$ be an integer. We regard X^d as the set $\{\varphi: [d] \rightarrow X\}$ of all maps from $[d]$ to X . We equip X^d with the product topology and the diagonal action of G given by $(g\varphi)(a) := g\varphi(a)$ for all $\varphi \in X^d$ and $a \in [d]$. Define the metrics ρ_2 and ρ_∞ on X^d by setting

$$\rho_2(\varphi, \psi) := \left(\frac{1}{d} \sum_{a \in [d]} (\rho(\varphi(a), \psi(a)))^2 \right)^{1/2} \quad \text{and}$$

$$\rho_\infty(\varphi, \psi) := \max_{a \in [d]} \rho(\varphi(a), \psi(a))$$

for all $\varphi, \psi \in X^d$.

Given a nonempty finite subset $F \subset G$, a map $\sigma: G \rightarrow \text{Sym}(d)$, and a real number $\delta > 0$, we denote by $\text{Map}(X, \rho, F, \delta, \sigma)$ the set consisting of all maps $\varphi: [d] \rightarrow X$ such that

$$\rho_2(\varphi \circ \sigma_s, s\varphi) < \delta \quad \text{for all } s \in F,$$

equivalently,

$$(2.5) \quad \sum_{a \in [d]} \rho(\varphi(\sigma_s a), s\varphi(a))^2 < d\delta^2 \quad \text{for all } s \in F.$$

Given $\varepsilon > 0$, we denote by $N_\varepsilon(\text{Map}(X, \rho, F, \delta, \sigma), \rho_\infty)$ the maximal cardinality of a $(\rho_\infty, \varepsilon)$ -separated subset $Z \subset \text{Map}(X, \rho, F, \delta, \sigma)$. Note that $N_\varepsilon(\text{Map}(X, \rho, F, \delta, \sigma), \rho_\infty) = -\infty$ if $\text{Map}(X, \rho, F, \delta, \sigma) = \emptyset$ and otherwise it is finite since X^d is compact.

The *sofic topological entropy* of (X, G) with respect to Σ is then defined as

$$h_\Sigma(X, G) := \sup_{\varepsilon > 0} \inf_{F \in \mathcal{F}(G)} \inf_{\delta > 0} \limsup_{j \rightarrow \infty} \frac{1}{d_j} \log N_\varepsilon(\text{Map}(X, \rho, F, \delta, \sigma_j), \rho_\infty).$$

It can be shown that $h_\Sigma(X, G) \in \{-\infty\} \cup [0, \infty]$ does not depend on the choice of the metric ρ for (X, G) (see [26, Proposition 10.25]). It follows in particular that $h_\Sigma(X, G)$ is a *topological conjugacy invariant*, i.e., if (Y, G) is another dynamical system such that there exists a G -equivariant homeomorphism $f: X \rightarrow Y$ then $h_\Sigma(X, G) = h_\Sigma(Y, G)$.

The following is a generalization of a result by Keynes and Robertson [27] for expansive homeomorphisms (see also [19, Chapter 16] and, for a uniform version over amenable groups, [11, Corollary 4.21]).

Lemma 2.13. *Let G be a countable sofic group and let (X, G) be an expansive dynamical system. Fix a compatible metric ρ on X and let $c > 0$ be an expansivity constant for (X, G) relative to ρ . Let $\Sigma = (d_j, \sigma_j)_{j \in \mathbb{N}}$ be a sofic approximation for G . Then*

$$(2.6) \quad h_\Sigma(X, G) = \inf_{F \in \mathcal{F}(G)} \inf_{\delta > 0} \limsup_{j \rightarrow \infty} \frac{1}{d_j} \log N_{c/2}(\text{Map}(X, \rho, F, \delta, \sigma_j), \rho_\infty) < +\infty.$$

Proof. Clearly $h_\Sigma(X, G)$ is no less than the right hand side in (2.6). By virtue of [26, Proposition 10.23], it suffices to show that for any $\varepsilon > 0$ one has

$$\begin{aligned} \inf_{F \in \mathcal{F}(G)} \inf_{\delta > 0} \limsup_{j \rightarrow \infty} \frac{1}{d_j} \log N_\varepsilon(\text{Map}(X, \rho, F, \delta, \sigma_j), \rho_2) \\ \leq \inf_{F \in \mathcal{F}(G)} \inf_{\delta > 0} \limsup_{j \rightarrow \infty} \frac{1}{d_j} \log N_{c/2}(\text{Map}(X, \rho, F, \delta, \sigma_j), \rho_\infty). \end{aligned}$$

Let $\varepsilon > 0$. By Lemma 2.2 we can find $K \in \mathcal{F}(G)$ such that for any $x, y \in X$ with $\rho_K(x, y) \leq c$ one has $\rho(x, y) < \varepsilon/4$.

Let $F \in \mathcal{F}(G)$ containing K and let $\delta > 0$ be small enough so that $\delta < (c/4)^2$ and $2|K|\delta \text{diam}(X, \rho)^2 < (\varepsilon/4)^2$. It suffices to show that for any $d \in \mathbb{N}$ and any map $\sigma: G \rightarrow \text{Sym}(d)$ one has

$$N_\varepsilon(\text{Map}(X, \rho, F, \delta, \sigma), \rho_2) \leq N_{c/2}(\text{Map}(X, \rho, F, \delta, \sigma), \rho_\infty).$$

For each $\varphi \in \text{Map}(X, \rho, F, \delta, \sigma)$ set

$$W_\varphi := \{a \in [d] : \rho(\varphi(\sigma_s a), s\varphi(a)) < \delta^{1/2} \text{ for all } s \in K\}.$$

We claim that

$$(2.7) \quad \frac{|W_\varphi|}{d} \geq 1 - |K|\delta.$$

In order to prove the claim, for $s \in K$ let us set $W_{\varphi, s} := \{a \in [d] : \rho(\varphi(\sigma_s a), s\varphi(a)) < \delta^{1/2}\}$. By applying the Chebyshev inequality we have

$$\begin{aligned} |[d] \setminus W_{\varphi, s}| &= |\{a \in [d] : \rho(\varphi(\sigma_s a), s\varphi(a)) \geq \delta^{1/2}\}| \\ &\leq \frac{1}{\delta} \sum_{a \in [d] \setminus W_{\varphi, s}} \rho(\varphi(\sigma_s a), s\varphi(a))^2 \\ &\leq \frac{1}{\delta} \sum_{a \in [d]} \rho(\varphi(\sigma_s a), s\varphi(a))^2 \\ &\quad (\text{by (2.5)}) \leq d\delta. \end{aligned}$$

We deduce that

$$|[d] \setminus W_\varphi| = \left| \bigcup_{s \in K} ([d] \setminus W_{\varphi, s}) \right| \leq d|K|\delta,$$

and the claim follows.

Take a $(\rho_\infty, c/2)$ -separated subset Φ of $\text{Map}(X, \rho, F, \delta, \sigma)$ with

$$|\Phi| = N_{c/2}(\text{Map}(X, \rho, F, \delta, \sigma), \rho_\infty).$$

Let $\psi \in \text{Map}(X, \rho, F, \delta, \sigma)$. Then $\rho_\infty(\psi, \varphi) < c/2$ for some $\varphi \in \Phi$. For every $a \in W_\varphi \cap W_\psi$, we have

$$\begin{aligned} \rho(s\varphi(a), s\psi(a)) &\leq \rho(s\varphi(a), \varphi(\sigma_s a)) + \rho(\varphi(\sigma_s a), \psi(\sigma_s a)) + \rho(\psi(\sigma_s a), s\psi(a)) \\ &\leq \delta^{1/2} + c/2 + \delta^{1/2} \leq c \end{aligned}$$

for all $s \in K$, and hence $\rho(\varphi(a), \psi(a)) < \varepsilon/4$ by our choice of K . Therefore

$$\begin{aligned} \rho_2(\varphi, \psi) &\leq \left(\frac{|W_\varphi \cap W_\psi|}{d} \left(\frac{\varepsilon}{4}\right)^2 + \left(1 - \frac{|W_\varphi \cap W_\psi|}{d}\right) \text{diam}(X, \rho)^2 \right)^{1/2} \\ (\text{by (2.7)}) &\leq \left(\left(\frac{\varepsilon}{4}\right)^2 + 2|K|\delta \text{diam}(X, \rho)^2 \right)^{1/2} < \varepsilon/2. \end{aligned}$$

Then

$$N_\varepsilon(\text{Map}(X, \rho, F, \delta, \sigma), \rho_2) \leq |\Phi| = N_{c/2}(\text{Map}(X, \rho, F, \delta, \sigma), \rho_\infty),$$

as desired. \square

3. PROOFS

Proof of Proposition 1.2. Take a compatible metric ρ for X and let $c > 0$ be an expansivity constant for (X, G) relative to ρ . Take a point $x_0 \in X \setminus Y$ and set $\varepsilon := \min(\rho(x_0, Y), c)/6 > 0$. Take a maximal (ρ, ε) -separated subset Z of X .

Since (X, G) is expansive and has the strong TMP, by Remark 2.5 it has the uniform strong TMP. Thus we can find $\tau > 0$ and $F_1 \in \mathcal{F}(G)$ containing 1_G such that for any $A \in \mathcal{F}(G)$, any $F_1 A$ -separated $V \subset G$, any $x_s \in X$ for $s \in V$, and any $y \in X$ satisfying $\rho_{(F_1 A \setminus A)s}(x_s, y) \leq \tau$ for all $s \in V$, there is some $z \in X$ so that $\rho_{F_1 A s}(x_s, z) \leq \varepsilon$ for all $s \in V$ and $\rho_{G \setminus AV}(y, z) \leq \varepsilon$.

Moreover, by the weak specification property, we can find some symmetric $F_2 \in \mathcal{F}(G)$ containing 1_G such that for any $A_1, A_2 \subset G$ satisfying $F_2 A_1 \cap F_2 A_2 = \emptyset$ and any $x_1, x_2 \in X$ there is some $z \in X$ such that $\rho_{A_1}(x_1, z) \leq \min(\tau/2, \varepsilon)$ and $\rho_{A_2}(x_2, z) \leq \min(\tau/2, \varepsilon)$. In particular (by taking $A_1 := \{1_G\}$ and $A_2 := G \setminus F_2^2$), for each $y \in Y$, there is some $z_y \in X$ such that $\rho(x_0, z_y) \leq \min(\tau/2, \varepsilon)$ and $\rho_{G \setminus F_2^2}(y, z_y) \leq \min(\tau/2, \varepsilon)$.

Let now $K \in \mathcal{F}(G)$ and $\delta, \kappa > 0$.

By Stirling's approximation formula (cf. Lemma A.1), there is some $0 < \gamma < 1$ such that for any $d \in \mathbb{N}$, the number of subsets $D \subset [d]$ with $|D| \leq \gamma d$ is at most $e^{\kappa d}$. Take $0 < \theta < \min(1/4, \gamma)$ such that

$$(3.1) \quad |Z|^{2\theta} \leq e^\kappa$$

and $4\theta \text{diam}(X, \rho)^2 \leq (\delta/2)^2$.

Since c is an expansivity constant for (X, G) , by Lemma 2.2 we can find some $K_1 \in \mathcal{F}(G)$ containing 1_G such that for any $x_1, x_2 \in X$ with $\rho_{K_1}(x_1, x_2) \leq c$, one has $\rho(x_1, x_2) < \delta/2$.

By compactness of X , the continuous action of G is indeed uniformly continuous. As a consequence, we can find $\xi > 0$ such that

$$(3.2) \quad \rho(x, y) \leq \xi \Rightarrow \rho_{F_1 F_2^2}(x, y) \leq \tau/2 \quad \text{and} \quad \rho_{K_1}(x, y) \leq \varepsilon$$

for all $x, y \in X$.

Set $F := F_1 F_2^2 F_1^{-1} K_1(\{1_G\} \cup K) \in \mathcal{F}(G)$, and take $\zeta > 0$ such that

$$(3.3) \quad \zeta \leq \xi^2 \quad \text{and} \quad \zeta|F| \leq \theta.$$

Let $j_0 \in \mathbb{N}$ be large enough so that for all $j \geq j_0$, setting for simplicity $d := d_j$ and $\sigma := \sigma_j$, we have that (cf. Section 2.11)

$$(3.4) \quad \frac{|U|}{d} \geq 1 - \theta,$$

where U denotes the set of $a \in [d]$ satisfying that $\sigma_s a \neq \sigma_t a$ for all distinct $s, t \in F$ and $\sigma_{st} a = \sigma_s \sigma_t a$ for all $s, t \in F$. Note that $\sigma_{1_G} a = a$ for all $a \in U$, since $1_G \in F$.

For any $\varphi, \psi: [d] \rightarrow X$ and any $\Theta \subset [d]$, we put

$$\rho_{\infty, \Theta}(\varphi, \psi) := \max_{a \in \Theta} \rho(\varphi(a), \psi(a)).$$

As, by our assumptions, $h_\Sigma(X, G) \geq 0$, the statement would trivially hold if $h_\Sigma(Y, G) = -\infty$. Thus, we may assume that $\text{Map}(Y, \rho, F, \zeta, \sigma) \neq \emptyset$. For $\varphi \in \text{Map}(Y, \rho, F, \zeta, \sigma)$ we then set

$$W_\varphi := \{a \in [d] : \rho(\varphi(\sigma_s a), s\varphi(a)) \leq \zeta^{1/2} \text{ for all } s \in F\}.$$

As in the proof of Lemma 2.13, we have (cf. (2.7)) $\frac{|W_\varphi|}{d} \geq 1 - |F|\zeta$ so that as $|F|\zeta \leq \theta$ (cf. (3.3)),

$$(3.5) \quad \frac{|W_\varphi|}{d} \geq 1 - \theta.$$

Take a $(\rho_\infty, c/2)$ -separated subset Φ of $\text{Map}(Y, \rho, F, \zeta, \sigma)$ such that

$$(3.6) \quad |\Phi| = N_{c/2}(\text{Map}(Y, \rho, F, \zeta, \sigma), \rho_\infty).$$

By our choice of θ and γ (cf. (3.1)) there is some $\Phi_1 \subset \Phi$ such that

$$(3.7) \quad |\Phi_1| \geq |\Phi|e^{-\kappa d}$$

and the set W_φ is the same for all $\varphi \in \Phi_1$. Set $W := W_\varphi$ for $\varphi \in \Phi_1$, and $U' := U \cap W$. Then, keeping in mind (3.4) and (3.5), we have $\frac{|U'|}{d} \geq 1 - 2\theta$.

Since Z is a maximal (ρ, ε) -separated subset of X , for every $\varphi \in \Phi_1$ and $a \in [d] \setminus U'$ we can find an element $f_\varphi(a) \in Z$ such that $\rho(\varphi(a), f_\varphi(a)) < \varepsilon$. This defines a map $f_\varphi: [d] \setminus U' \rightarrow Z$. Then, there is a set $\Phi_2 \subset \Phi_1$ such that

$$|\Phi_2| \geq |\Phi_1| \cdot |Z|^{-|[d] \setminus U'|} \geq |\Phi_1| \cdot |Z|^{-2\theta d} \geq |\Phi_1|e^{-\kappa d}$$

and the map f_φ is the same for all $\varphi \in \Phi_2$. Note that by (3.7) we have

$$(3.8) \quad |\Phi_2| \geq |\Phi|e^{-2\kappa d}.$$

Now, for any distinct $\varphi, \psi \in \Phi_2$, we have $\rho_{\infty, [d] \setminus U'}(\varphi, \psi) < 2\varepsilon \leq c/2$ and $\rho_\infty(\varphi, \psi) \geq c/2$, and hence

$$(3.9) \quad \rho_{\infty, U'}(\varphi, \psi) \geq c/2.$$

Take a maximal subset Λ of U' subject to the condition that $\sigma(F_1 F_2^2)w_1 \cap \sigma(F_1 F_2^2)w_2 = \emptyset$ for all distinct $w_1, w_2 \in \Lambda$. Then $\sigma(F_1 F_2^2)^{-1} \sigma(F_1 F_2^2) \Lambda \supseteq U'$, and hence (recalling that

$$0 < \theta \leq 1/4)$$

$$(3.10) \quad \frac{|\Lambda|}{d} \geq \frac{|U'|}{|F_1 F_2^2|^2 d} \geq \frac{1 - 2\theta}{|F_1 F_2^2|^2} \geq \frac{1}{2|F_1 F_2^2|^2}.$$

Let $B \subset \Lambda$. Again, since Z is a maximal (ρ, ε) -separated subset of X , for every $\varphi \in \Phi_2$ and $a \in \sigma(F_1 F_2^2)B$ we can find an element $f_{B,\varphi}(a) \in Z$ such that $\rho(\varphi(a), f_{B,\varphi}(a)) < \varepsilon$. This defines a map $f_{B,\varphi}: \sigma(F_1 F_2^2)B \rightarrow Z$. Then there is a subset $\Phi_B \subset \Phi_2$ such that

$$|\Phi_B| \geq |\Phi_2| \cdot |Z|^{-|\sigma(F_1 F_2^2)B|} = |\Phi_2| \cdot |Z|^{-|F_1 F_2^2| \cdot |B|}$$

and the map $f_{B,\varphi}$ is the same for all $\varphi \in \Phi_B$. Note that by (3.8) we have

$$(3.11) \quad |\Phi_B| \geq |\Phi| e^{-2\kappa d} |Z|^{-|F_1 F_2^2| \cdot |B|}.$$

For any distinct $\varphi, \psi \in \Phi_B$, we have $\rho_{\infty, \sigma(F_1 F_2^2)B}(\varphi, \psi) < 2\varepsilon \leq c/2$ and $\rho_{\infty, U'}(\varphi, \psi) \geq c/2$, and hence

$$(3.12) \quad \rho_{\infty, U' \setminus \sigma(F_1 F_2^2)B}(\varphi, \psi) \geq c/2.$$

Let $a \in U'$. Denote by V_a the set of all $s \in F$ such that $\sigma_s a \in \Lambda$. For any $s \in V_a$, since $F_1 F_2^2 \subset F$ and $a \in U$, we have $\sigma(F_1 F_2^2 s)a = \sigma(F_1 F_2^2)\sigma_s a$. Since $a \in U$, for any distinct s, t in V_a , we have $\sigma_s a \neq \sigma_t a$, and hence

$$\sigma(F_1 F_2^2 s)a \cap \sigma(F_1 F_2^2 t)a = \sigma(F_1 F_2^2)\sigma_s a \cap \sigma(F_1 F_2^2)\sigma_t a = \emptyset.$$

It follows that V_a is $F_1 F_2^2$ -separated.

With a subset $B \subset \Lambda$ and $\varphi \in \Phi_B$ we associate a map $\bar{\varphi}_B: [d] \rightarrow X$ defined as follows.

Let $a \in [d]$. Suppose first that $a \in U'$. Denote by $V_{B,a}$ the set of all $s \in F$ such that $\sigma_s a \in B$. Then $V_{B,a} \subset V_a$ and hence $V_{B,a}$ is $F_1 F_2^2$ -separated. Let $s \in V_{B,a}$ and consider $\varphi(\sigma_s a) \in Y$. By our choice of F_2 , there is a point $z_{\varphi(\sigma_s a)} \in X$ with $\rho(x_0, z_{\varphi(\sigma_s a)}) \leq \min(\tau/2, \varepsilon)$ and $\rho_{G \setminus F_2^2}(z_{\varphi(\sigma_s a)}, \varphi(\sigma_s a)) \leq \min(\tau/2, \varepsilon)$. By our choice of ξ , since $\rho(\varphi(\sigma_s a), s\varphi(a)) \leq \xi^{1/2} \leq \xi$, we have $\rho_{F_1 F_2^2}(\varphi(\sigma_s a), s\varphi(a)) \leq \tau/2$. Setting $x_{\varphi,a,s} := s^{-1}z_{\varphi(\sigma_s a)} \in X$, we have

$$\begin{aligned} \rho_{(F_1 F_2^2 \setminus F_2^2)s}(x_{\varphi,a,s}, \varphi(a)) &\leq \rho_{(F_1 F_2^2 \setminus F_2^2)s}(s^{-1}z_{\varphi(\sigma_s a)}, s^{-1}\varphi(\sigma_s a)) + \rho_{(F_1 F_2^2 \setminus F_2^2)s}(s^{-1}\varphi(\sigma_s a), \varphi(a)) \\ &= \rho_{F_1 F_2^2 \setminus F_2^2}(z_{\varphi(\sigma_s a)}, \varphi(\sigma_s a)) + \rho_{F_1 F_2^2 \setminus F_2^2}(\varphi(\sigma_s a), s\varphi(a)) \\ &\leq \tau/2 + \tau/2 = \tau. \end{aligned}$$

By our choice of F_1 and τ (coming from the uniform sTMP of (X, G)), there is some $\bar{\varphi}_B(a) \in X$ such that

$$(3.13) \quad \rho_{F_1 F_2^2 s}(x_{\varphi,a,s}, \bar{\varphi}_B(a)) \leq \varepsilon \quad \text{for all } s \in V_{B,a}$$

and

$$(3.14) \quad \rho_{G \setminus F_2^2 V_{B,a}}(\varphi(a), \bar{\varphi}_B(a)) \leq \varepsilon.$$

For $a \in [d] \setminus U'$, we set $\bar{\varphi}_B(a) := x_0$.

Claim 1. $\bar{\varphi}_B \in \text{Map}(X, \rho, K, \delta, \sigma)$.

Proof of Claim 1. Let $t \in K$, let $a \in U' \cap \sigma_t^{-1}U'$, and let $g \in K_1$.

Suppose $gt \in F_1F_2^2V_{B,a}$, say $gt = hs$ with $h \in F_1F_2^2$ and $s \in V_{B,a}$. Then $h^{-1}g \in F_2^2F_1^{-1}K_1 \subset F$ and $\sigma_{h^{-1}g}\sigma_ta = \sigma_{h^{-1}gt}a = \sigma_sa \in B$, and hence $h^{-1}g \in V_{B,\sigma_ta}$. In this case we have

$$(3.15) \quad gx_{\varphi,\sigma_ta,h^{-1}g} = g(h^{-1}g)^{-1}z_{\varphi(\sigma_{h^{-1}g}\sigma_ta)} = hz_{\varphi(\sigma_{h^{-1}g}\sigma_ta)} = gts^{-1}z_{\varphi(\sigma_sa)} = gtx_{\varphi,a,s},$$

and hence using $g = h(h^{-1}g) \in F_1F_2^2V_{B,\sigma_ta}$, we have

$$\begin{aligned} \rho(g\bar{\varphi}_B(\sigma_ta), gt\bar{\varphi}_B(a)) &\leq \rho(g\bar{\varphi}_B(\sigma_ta), gx_{\varphi,\sigma_ta,h^{-1}g}) + \rho(gx_{\varphi,\sigma_ta,h^{-1}g}, gtx_{\varphi,a,s}) \\ &\quad + \rho(gtx_{\varphi,a,s}, gt\bar{\varphi}_B(a)) \\ &\quad (\text{by (3.15)}) = \rho(g\bar{\varphi}_B(\sigma_ta), gx_{\varphi,\sigma_ta,h^{-1}g}) + \rho(gtx_{\varphi,a,s}, gt\bar{\varphi}_B(a)) \\ &\quad (\text{as } gt = hs) = \rho(g\bar{\varphi}_B(\sigma_ta), gx_{\varphi,\sigma_ta,h^{-1}g}) + \rho(hsx_{\varphi,a,s}, hs\bar{\varphi}_B(a)) \\ &\quad (\text{by (3.13)}) \leq \varepsilon + \varepsilon \leq c. \end{aligned}$$

Suppose $g \in F_1F_2^2V_{B,\sigma_ta}$, say $g = h_1s_1$ with $h_1 \in F_1F_2^2$ and $s_1 \in V_{B,\sigma_ta}$. Then $h_1^{-1}gt \in F_2^2F_1^{-1}K_1K \subset F$ and $\sigma_{h_1^{-1}gt}a = \sigma_{h_1^{-1}g}\sigma_ta = \sigma_{s_1}\sigma_ta \in B$, and hence $h_1^{-1}gt \in V_{B,a}$. In this case we have $gt = h_1(h_1^{-1}gt) \in F_1F_2^2V_{B,a}$, and hence, by the above, we get $\rho(g\bar{\varphi}_B(\sigma_ta), gt\bar{\varphi}_B(a)) \leq c$.

Finally, suppose that $gt \notin F_1F_2^2V_{B,a}$ and $g \notin F_1F_2^2V_{B,\sigma_ta}$. Since $\rho(\varphi(\sigma_ta), t\varphi(a)) \leq \zeta^{1/2} \leq \xi$, and recalling that $g \in K_1$, we have $\rho(g\varphi(\sigma_ta), gt\varphi(a)) \leq \varepsilon$ by (3.2). As a consequence,

$$\begin{aligned} \rho(g\bar{\varphi}_B(\sigma_ta), gt\bar{\varphi}_B(a)) &\leq \rho(g\bar{\varphi}_B(\sigma_ta), g\varphi(\sigma_ta)) + \rho(g\varphi(\sigma_ta), gt\varphi(a)) + \rho(gt\varphi(a), gt\bar{\varphi}_B(a)) \\ &\quad (\text{by (3.14)}) \leq \varepsilon + \varepsilon + \varepsilon \leq c. \end{aligned}$$

We conclude that $\rho_{K_1}(\bar{\varphi}_B(\sigma_ta), t\bar{\varphi}_B(a)) \leq c$, and hence $\rho(\bar{\varphi}_B(\sigma_ta), t\bar{\varphi}_B(a)) < \delta/2$ by our choice of K_1 . Now we get

$$\begin{aligned} \rho_2(\bar{\varphi}_B \circ \sigma_t, t\bar{\varphi}_B) &\leq \left((\delta/2)^2 + \left(1 - \frac{|U' \cap \sigma_t^{-1}U'|}{d}\right) \text{diam}(X, \rho)^2 \right)^{1/2} \\ &\leq ((\delta/2)^2 + 4\theta \text{diam}(X, \rho)^2)^{1/2} \\ &\leq \delta. \end{aligned}$$

This proves our claim. □

Let $B \subset \Lambda$ and $\varphi \in \Phi_B$. For any $a \in B$, we have $1_G \in V_{B,a}$, and hence

$$(3.16) \quad \rho(\bar{\varphi}_B(a), x_0) \leq \rho(\bar{\varphi}_B(a), x_{\varphi,a,1_G}) + \rho(x_{\varphi,a,1_G}, x_0) \leq \varepsilon + \rho(z_{\varphi(a)}, x_0) \leq \varepsilon + \varepsilon = 2\varepsilon.$$

For any $a \in U' \setminus \sigma(F_1F_2^2)B$, we have $1_G \notin F_1F_2^2V_{B,a}$, and hence

$$(3.17) \quad \rho(\bar{\varphi}_B(a), \varphi(a)) \leq \varepsilon.$$

In particular, (3.17) holds for all $a \in \Lambda \setminus B \subset U' \setminus \sigma(F_1F_2^2)B$.

Let us set $\bar{\Phi} := \{\bar{\varphi}_B : B \subset \Lambda \text{ and } \varphi \in \Phi_B\} \subset \text{Map}(X, \rho, K, \delta, \sigma)$.

Claim 2. $\bar{\Phi}$ is $(\rho_\infty, \varepsilon)$ -separated.

Proof of Claim 2. For any distinct $B_1, B_2 \subset \Lambda$ and $\varphi \in \Phi_{B_1}, \psi \in \Phi_{B_2}$, we have for, say, $a \in B_1 \setminus B_2$,

$$\begin{aligned} \rho_\infty(\bar{\varphi}_{B_1}, \bar{\psi}_{B_2}) &\geq \rho(\bar{\varphi}_{B_1}(a), \bar{\psi}_{B_2}(a)) \\ &\geq \rho(x_0, \psi(a)) - \rho(\bar{\varphi}_{B_1}(a), x_0) - \rho(\bar{\psi}_{B_2}(a), \psi(a)) \\ (\text{by (3.16), (3.17), and } \varepsilon \leq \rho(x_0, Y)/6) &\geq 4\varepsilon - 2\varepsilon - \varepsilon = \varepsilon. \end{aligned}$$

For any $B \subset \Lambda$ and any distinct $\varphi, \psi \in \Phi_B$, by (3.12) we have $\rho(\varphi(a), \psi(a)) \geq c/2$ for some $a \in U' \setminus \sigma(F_1 F_2^2)B$, and hence

$$\begin{aligned} \rho_\infty(\bar{\varphi}_B, \bar{\psi}_B) &\geq \rho(\bar{\varphi}_B(a), \bar{\psi}_B(a)) \\ &\geq \rho(\varphi(a), \psi(a)) - \rho(\varphi(a), \bar{\varphi}_B(a)) - \rho(\psi(a), \bar{\psi}_B(a)) \\ (\text{by (3.17)}) &\geq c/2 - \varepsilon - \varepsilon \geq \varepsilon. \end{aligned}$$

This proves our claim. \square

We need the following elementary fact. Given a finite set Λ and a variable t , we have

$$(3.18) \quad \sum_{B \subset \Lambda} t^{|B|} = \sum_{j=0}^{|\Lambda|} \binom{|\Lambda|}{j} t^j = (1+t)^{|\Lambda|}.$$

We are now in a position to prove the entropic inequality and conclude the proof of the proposition.

$$\begin{aligned} N_\varepsilon(\text{Map}(X, \rho, K, \delta, \sigma), \rho_\infty) &\geq |\bar{\Phi}| \geq \sum_{B \subset \Lambda} |\Phi_B| \\ (\text{by (3.11)}) &\geq \sum_{B \subset \Lambda} |\Phi| e^{-2\kappa d} |Z|^{-|F_1 F_2^2| \cdot |B|} \\ (\text{by (3.6) and (3.18)}) &= N_{c/2}(\text{Map}(Y, \rho, F, \zeta, \sigma), \rho_\infty) e^{-2\kappa d} (1 + |Z|^{-|F_1 F_2^2|})^{|\Lambda|} \\ (\text{by (3.10)}) &\geq N_{c/2}(\text{Map}(Y, \rho, F, \zeta, \sigma), \rho_\infty) e^{-2\kappa d} (1 + |Z|^{-|F_1 F_2^2|})^{d/(2|F_1 F_2^2|^2)}. \end{aligned}$$

It follows that

$$\begin{aligned} &\limsup_{i \rightarrow \infty} \frac{1}{d_i} \log N_\varepsilon(\text{Map}(X, \rho, K, \delta, \sigma_i), \rho_\infty) \\ &\geq \limsup_{i \rightarrow \infty} \frac{1}{d_i} \log N_{c/2}(\text{Map}(Y, \rho, F, \zeta, \sigma_i), \rho_\infty) - 2\kappa + \frac{1}{2|F_1 F_2^2|^2} \log(1 + |Z|^{-|F_1 F_2^2|}) \\ &\geq h_\Sigma(Y, G) - 2\kappa + \frac{1}{2|F_1 F_2^2|^2} \log(1 + |Z|^{-|F_1 F_2^2|}), \end{aligned}$$

where in the 2nd inequality we apply Lemma 2.13 to (Y, G) . Letting $\kappa \rightarrow 0$, we obtain

$$\limsup_{i \rightarrow \infty} \frac{1}{d_i} \log N_\varepsilon(\text{Map}(X, \rho, K, \delta, \sigma_i), \rho_\infty) \geq h_\Sigma(Y, G) + \frac{1}{2|F_1 F_2^2|^2} \log(1 + |Z|^{-|F_1 F_2^2|}).$$

Taking the infimum over K and δ , we get

$$h_\Sigma(X, G) \geq h_\Sigma(Y, G) + \frac{1}{2|F_1 F_2|^2} \log(1 + |Z|^{-|F_1 F_2^2|}) > h_\Sigma(Y, G).$$

□

Proof of Theorem 1.1. Let $f: X \rightarrow X$ be an injective endomorphism of the dynamical system (X, G) . By continuity of f and since X is compact and Hausdorff, f yields a homeomorphism between X and $Y := f(X)$. Moreover, since f is G -equivariant, $Y \subset X$ is G -invariant. It follows that f in fact establishes a conjugacy between the dynamical systems (X, G) and (Y, G) so that $h_\Sigma(X, G) = h_\Sigma(Y, G)$. By Proposition 1.2, Y cannot be a proper subset of X . Thus $f(X) = Y = X$, that is, f is surjective. This shows that (X, G) is surjunctive. □

Proof of Corollary 1.4. Deninger and Schmidt [18, Theorem 3.2] proved that a principal algebraic dynamical system (X_f, G) is expansive if and only if f is invertible in $\ell^1(G)$ and Ren [33, Theorem 1.2] proved that every expansive principal algebraic action has the weak specification property. The proofs of the analogous statements for (X_A, G) , $A \in \text{Mat}_n(\mathbb{Z}[G])$, $n \geq 1$, follow essentially the same lines: the generalization of the Deninger-Schmidt result to finitely-presented algebraic actions is straightforward (see [16, Theorem 3.1] for an even more general setting). The generalization of Ren's result goes as follows. First of all, recall that, for subshifts, the weak specification property and strong irreducibility coincide (cf. Proposition 2.10 and Remark 2.11) and observe that for any nonempty finite set B , the full shift (B^G, G) is strongly irreducible and whence it has the weak specification property. Secondly, it is easy to see that weak specification property passes to factors, that is, if (X, G) and (Y, G) are two dynamical systems such that (X, G) satisfies the weak specification property and there exists a G -equivariant continuous surjective map $f: X \rightarrow Y$, then (Y, G) satisfies the weak specification property as well, as observed in [29, Section 2.1] (see also [11, Proposition 6.2] and [12, Exercise 1.66]). Suppose that $A \in \text{Mat}_n(\mathbb{Z}[G])$ is invertible in $\text{Mat}_n(\ell^1(G))$. Set $\|A\|_1 := \sum_{j,k} \|A_{j,k}\|_1$, where, $\|A_{j,k}\|_1 := \sum_{g \in G} |A_{j,k}(g)|$ denotes the ℓ^1 -norm of the (j, k) th entry $A_{j,k} \in \ell^1(G)$ of the matrix A . By virtue of (the matrix version of) [18, Proposition 4.2] and [33, Lemma 4.1], one has that (X_A, G) is a factor of the full shift (B^G, G) , where $B := \{(b_1, b_2, \dots, b_n) \in \mathbb{Z}^n : |b_j| \leq \frac{1}{2}\|A\|_1, 1 \leq j \leq n\}$. As a consequence, (X_A, G) has the weak specification property. Moreover, Meyerovitch [31, Theorem 3.4] proved that the finitely-presented dynamical system (X_A, G) associated with an element $A \in \text{Mat}_n(\mathbb{Z}[G])$ which is invertible in $\text{Mat}_n(\ell^1(G))$ has the pseudo-orbit tracing property (POTP). Finally, every dynamical system with the POTP has the strong topological Markov property (sTMP) [2, Proposition 3.10]. In conclusion, (X_A, G) is expansive and satisfies the weak specification property and the strong topological Markov property. The statement then follows from Theorem 1.1. □

Proof of Theorem 1.5. It is a result of Deninger that for any action of a countably infinite amenable group G on a compact metrizable group X by continuous automorphisms, the topological entropy is equal to the measure entropy $h_{\mu_X}(X, G)$ of the normalized Haar measure μ_X [17, Theorem 2.2] (though Deninger assumed X to be abelian, the proof there

does not use this assumption). It was shown in [16, Corollaries 7.5 and 8.4] that such an action has CPE if and only if it has CPE with respect to μ_X , i.e., every nontrivial factor of the probability-measure-preserving action $G \curvearrowright (X, \mu_X)$ has positive entropy. Furthermore, it was shown in [16, Theorem 8.6] that such an action with finite measure entropy of μ_X has CPE with respect to μ_X if and only if, for every G -invariant Borel probability measure ν of X not equal to μ_X , one has $h_\nu(X, G) < h_{\mu_X}(X, G)$.

Let now $f: X \rightarrow X$ be an injective endomorphism of (X, G) . We have $h_{f_*\mu_X}(X, G) = h_{f_*\mu_X}(f(X), G) = h_{\mu_X}(X, G)$ so that $f_*\mu_X = \mu_X$. Since μ_X has full support, we deduce that $f(X) = X$, that is, f is surjective. This shows that the dynamical system (X, G) is surjunctive. \square

4. EXAMPLES

Example 4.1 (The hard-ball model). Take $G = \mathbb{Z}$ and $A = \{0, 1\}$. Then the subset $X \subset A^G$, consisting of all $x: \mathbb{Z} \rightarrow \{0, 1\}$ such that $(x(n), x(n+1)) \neq (1, 1)$ for all $n \in \mathbb{Z}$, is a subshift. This subshift is called the *golden mean subshift*.

More generally, take $G = \mathbb{Z}^d$ and $A = \{0, 1\}$. Then the subset $X \subset A^G$, consisting of all $x \in A^G$ such that $(x(g), x(g + e_i)) \neq (1, 1)$ for all $g \in G$, where $(e_i)_{1 \leq i \leq d}$ is the canonical basis of \mathbb{Z}^d , is a subshift. This subshift is called the *hard-ball model*.

The hard-ball model is strongly irreducible and of finite type (and therefore splicable). To see this it suffices to convince ourselves that the set $S := \{g = (g_1, \dots, g_d) \in G : |g_i| \leq 1, i = 1, \dots, d\}$ witnesses both strong irreducibility and the finite type condition.

In [2, Example 6.6] the following generalization is presented. Let G be a group, let $F \subset G$ be a finite nonempty subset not containing 1_G . The F -hard-ball model of G is the subshift $X_{\text{hard}, F} \subset \{0, 1\}^G$ defined by setting

$$X_{\text{hard}, F} := \{x \in \{0, 1\}^G : x(g)x(gs) = 0 \text{ for all } g \in G \text{ and } s \in F\}.$$

Then $X_{\text{hard}, F}$ is a strongly irreducible subshift of finite type, and therefore has the POTP and thus the sTMP, equivalently the usTMP (cf. [2, Proposition 3.11] and Proposition 2.6). When G is sofic, given any sofic approximation Σ for G , one has $h_\Sigma(X_{\text{hard}, F}, G) \geq 0$, since the action has fixed points (in fact, only one, namely the constant 0-configuration).

We deduce from Theorem 1.1 that if G is sofic then the F -hard-ball model $(X_{\text{hard}, F}, G)$ is surjunctive.

Example 4.2 (Weiss' example). Benjy Weiss [36] found the following simple example of a shift of finite type that is not surjunctive. Let $A = \{0, 1, 2\}$ and $G = \mathbb{Z}$. Consider the subshift $X \subset A^G$ of finite type with defining set of admissible patterns $\mathcal{A} = \{00, 01, 11, 12, 22\} \subset A^2 = A^\Omega$, where $\Omega = \{0, 1\} \subset G$ (cf. Section 2.3). Thus any point x in X has at most one block of 1's, which, if finite is bordered by an infinite string of 0's to the left and 2's to the right. Set $S := \{-1, 0\} \subset G$ and define $\mu: A^S = A^2 \rightarrow A$ by setting

$$\mu(s, t) = \begin{cases} t & \text{if } (s, t) \neq (1, 2) \\ 1 & \text{if } (s, t) = (1, 2). \end{cases}$$

Then the associated cellular automaton $f: A^G \rightarrow A^G$ leaves X invariant (and therefore is an endomorphism of (X, G)) and works as follows: if the finite block of 1's occurs in $x \in X$, then in $f(x)$ the same block is elongated by one extra 1 on the right. Thus f is injective. Clearly, f is not surjective since a configuration with a single 1 will be in $X \setminus f(X)$. As a consequence (X, G) is not surjunctive. Note that (X, G) has the sTMP (indeed, the subshift X is of finite type) but it does not satisfy the weak specification property (indeed, the subshift X is not strongly irreducible).

APPENDIX A. STIRLING'S APPROXIMATION FORMULA

The following old result was widely used, without proof, in [25]. For the reader's convenience and the sake of completeness, we include its detailed proof.

Given a real number α we denote by $\lfloor \alpha \rfloor \in \mathbb{Z}$ the greatest integer less than or equal to α .

Lemma A.1. *Let $0 < \gamma < 1/2$. Then there exists $\kappa = \kappa(\gamma) > 0$ and $d_0 = d_0(\gamma) \in \mathbb{N}$ with*

$$(A.1) \quad \lim_{\gamma \rightarrow 0} \kappa(\gamma) = 0$$

such that

$$(A.2) \quad \sum_{j=0}^{\lfloor \gamma d \rfloor} \binom{d}{j} \leq e^{\kappa d}$$

for all $d \in \mathbb{N}$ such that $d \geq d_0$.

Proof. Let $d \geq 2/\gamma$ so that, on the one hand

$$(A.3) \quad \lfloor \gamma d + 1 \rfloor = \lfloor \gamma d \rfloor + 1 \leq 2\lfloor \gamma d \rfloor,$$

and, on the other hand, since $\gamma < 1/2$, one has $d \geq 4$ and therefore $d - \gamma d \geq 2$.

As $\gamma < 1/2$, we clearly have

$$\sum_{j=0}^{\lfloor \gamma d \rfloor} \binom{d}{j} \leq (\lfloor \gamma d \rfloor + 1) \binom{d}{\lfloor \gamma d \rfloor},$$

so that, by virtue of (A.3), we only need to find $\kappa = \kappa(\gamma)$ satisfying (A.1) and such that

$$(A.4) \quad 2\lfloor \gamma d \rfloor \binom{d}{\lfloor \gamma d \rfloor} \leq e^{\kappa d}.$$

Setting $\kappa := -2(\gamma \ln \gamma + (1 - \gamma) \ln(1 - \gamma)) > 0$, we clearly have that (A.1) holds. Consider the function $G(x) = e^{\kappa x/2} - x^3$. It is easy to see that there exists an integer $d_0 \geq 2/\gamma$ such that $G(x) > 0$ for all $x \geq d_0$.

In [26, Lemma 10.1] the following version of Stirling's classical formula is proved by purely elementary means:

$$(A.5) \quad e \left(\frac{m}{e} \right)^m \leq m! \leq em \left(\frac{m}{e} \right)^m$$

for all $m \in \mathbb{N}$ such that $m \geq 1$.

Let $d \geq d_0$. Then, setting $m := \lfloor \gamma d \rfloor$, we deduce that

$$\begin{aligned}
2\lfloor \gamma d \rfloor \binom{d}{\lfloor \gamma d \rfloor} &= 2m \binom{d}{m} = 2m \cdot \frac{d!}{m!(d-m)!} \\
&\leq 2m \cdot \frac{ed \left(\frac{d}{e}\right)^d}{e \left(\frac{m}{e}\right)^m \cdot e \left(\frac{d-m}{e}\right)^{d-m}} \\
&= 2m \cdot \frac{d}{e} \cdot d^d \cdot \frac{1}{m^m} \cdot \frac{1}{(d-m)^{d-m}} \\
(\text{since } \gamma d - 1 \leq m \leq \gamma d) &\leq 2\gamma d \cdot \frac{d}{e} \cdot d^d \cdot \frac{1}{(\gamma d - 1)^{\gamma d - 1}} \cdot \frac{1}{((1-\gamma)d)^{(1-\gamma)d}} \\
&= 2\gamma d \cdot \frac{d}{e} \cdot d^d \cdot \gamma d \cdot \left(\frac{\gamma d}{\gamma d - 1}\right)^{\gamma d - 1} \\
&\quad \cdot \frac{1}{d^d} \cdot \frac{1}{\gamma^{\gamma d}} \cdot \frac{1}{(1-\gamma)^{(1-\gamma)d}} \\
(\text{since } (\gamma d / (\gamma d - 1))^{\gamma d - 1} \leq e) &\leq 2d^3 \gamma^2 \cdot \frac{1}{\gamma^{\gamma d}} \cdot \frac{1}{(1-\gamma)^{(1-\gamma)d}} \\
(\text{since } \gamma < 1/2) &< d^3 \cdot \left(\frac{1}{\gamma^\gamma} \cdot \frac{1}{(1-\gamma)^{(1-\gamma)}}\right)^d \\
&= d^3 \cdot e^{-(\gamma \ln \gamma + (1-\gamma) \ln(1-\gamma))d} \\
(\text{as } d \geq d_0) &\leq e^{\kappa d}.
\end{aligned}$$

This proves (A.4) and the proof is complete. \square

REFERENCES

- [1] M. Achigar, A note on Anosov homeomorphisms, *Axioms* **8** (2019), no. 2, 54.
- [2] S. Barbieri, F. García-Ramos, and H. Li, Markovian properties of continuous group actions: algebraic actions, entropy and the homoclinic group, *Adv. Math.* **397** (2022), Paper No. 108196, 52 pp.
- [3] S. Barbieri, R. Gómez, B. Marcus, and S. Taati, Equivalence of relative Gibbs and relative equilibrium measures for actions of countable amenable groups, *Nonlinearity* **33** (2020), no. 5, 2409–2454.
- [4] S. Bhattacharya, T. Ceccherini-Silberstein, and M. Coornaert, Surjunctivity and topological rigidity of algebraic dynamical systems, *Ergodic Theory Dynam. Systems* **39** (2019), no. 3, 604–619.
- [5] R. Bowen, Periodic points and measures for Axiom A diffeomorphisms, *Trans. Amer. Math. Soc.* **154** (1971), 377–397.
- [6] R. Bowen, *On Axiom A diffeomorphisms*, American Mathematical Society, Providence, R.I., 1978. Regional Conference Series in Mathematics, No. 35.
- [7] B.F. Bryant, Expansive self-homeomorphisms of a compact metric space, *Amer. Math. Monthly* **69** (1962), 386–391.
- [8] V. Capraro and M. Lupini, *Introduction to sofic and hyperlinear groups and Connes’ embedding conjecture. With an appendix by Vladimir Pestov*, Lecture Notes in Mathematics, 2136. Springer, Cham, 2015.
- [9] T. Ceccherini-Silberstein and M. Coornaert, The Myhill property for strongly irreducible subshifts over amenable groups, *Monatsh. Math.* **165** (2012), 155–172.

- [10] T. Ceccherini-Silberstein and M. Coornaert, *Cellular automata and groups*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2010.
- [11] T. Ceccherini-Silberstein and M. Coornaert, Expansive actions with specification on uniform spaces, topological entropy, and the Myhill property, *J. Dyn. Control Syst.* **27** (2021), no. 3, 427–456.
- [12] T. Ceccherini-Silberstein and M. Coornaert, *Exercises in cellular automata and groups*, Springer Monographs in Mathematics, Springer Cham, 2023.
- [13] T. Ceccherini-Silberstein, F. Fiorenzi, and F. Scarabotti, The Garden of Eden theorem for cellular automata and for symbolic dynamical systems, in *Random walks and geometry*, Walter de Gruyter, Berlin, 2004, pp. 73–108.
- [14] T. Ceccherini-Silberstein, A. Machì, and F. Scarabotti, Amenable groups and cellular automata, *Ann. Inst. Fourier (Grenoble)* **49** (1999), 673–685.
- [15] N.-P. Chung and K. Lee, Topological stability and pseudo-orbit tracing property of group actions, *Proc. Amer. Math. Soc.* **146** (2018), no. 3, 1047–1057.
- [16] N.-P. Chung and H. Li, Homoclinic groups, IE groups, and expansive algebraic actions, *Invent. Math.* **199** (2015), no. 3, 805–858.
- [17] C. Deninger, Fuglede-Kadison determinants and entropy for actions of discrete amenable groups, *J. Amer. Math. Soc.* **19** (2006), no. 3, 737–758.
- [18] C. Deninger and K. Schmidt, Expansive algebraic actions of discrete residually finite amenable groups and their entropy, *Ergodic Theory Dynam. Systems* **27** (2007), no. 3, 769–786.
- [19] M. Denker, C. Grillenberger, and K. Sigmund, *Ergodic theory on compact spaces*, Lecture Notes in Mathematics, Vol. 527. Springer-Verlag, Berlin-New York, 1976.
- [20] M. Doucha, Garden of Eden and weakly periodic points for certain expansive actions of groups, *Ergodic Theory Dynam. Systems*, 1–22. doi:10.1017/etds.2022.37
- [21] F. Fiorenzi, Cellular automata and strongly irreducible shifts of finite type, *Theoret. Comput. Sci.* **299** (2003), 477–493.
- [22] W.H. Gottschalk, Some general dynamical systems, Recent advances in topological dynamics, pp. 120–125, Lecture Notes in Mathematics Vol. 318, Springer, Berlin, 1973.
- [23] M. Gromov, Endomorphisms of symbolic algebraic varieties, *J. Eur. Math. Soc. (JEMS)* **1** (1999), 109–197.
- [24] D. Kerr and H. Li, Entropy and the variational principle for actions of sofic groups, *Invent. Math.* **186** (2011), no. 3, 501–558.
- [25] D. Kerr and H. Li, Soficity, amenability, and dynamical entropy, *Amer. J. Math.* **135** (2013), no. 3, 721–761.
- [26] D. Kerr and H. Li, *Ergodic theory. Independence and dichotomies*. Springer Monographs in Mathematics. Springer, Cham, 2016.
- [27] H.B. Keynes and J.B. Robertson, Generators for topological entropy and expansiveness, *Math. Systems Theory* **3** (1969), 51–59.
- [28] W. Lawton, Note on symbolic transformation groups, *Notices of the AMS* **19** (1972), pA–376 (abstract).
- [29] H. Li, Garden of Eden and specification, *Ergodic Theory Dynam. Systems* **39** (2019), no. 11, 3075–3088.
- [30] D. Lind and K. Schmidt, Homoclinic points of algebraic \mathbf{Z}^d -actions, *J. Amer. Math. Soc.* **12** (1999), 953–980.
- [31] T. Meyerovitch, Pseudo-orbit tracing and algebraic actions of countable amenable groups, *Ergodic Theory Dynam. Systems* **39** (2019), no. 9, 2570–2591.
- [32] S. A. Morris, *Pontryagin duality and the structure of locally compact Abelian groups*, Cambridge University Press, Cambridge-New York-Melbourne, 1977. London Mathematical Society Lecture Note Series, No. 29.
- [33] X. Ren, Periodic measures are dense in invariant measures for residually finite amenable group actions with specification, *Discrete Contin. Dyn. Syst.* **38** (2018), no. 4, 1657–1667.

- [34] D. Ruelle, Statistical mechanics on a compact set with Z^ν action satisfying expansiveness and specification, *Trans. Amer. Math. Soc.* **187** (1973), 237–251.
- [35] P. Walters, On the pseudo-orbit tracing property and its relationship to stability. In The structure of attractors in dynamical systems (Proc. Conf., North Dakota State Univ., Fargo, N.D., 1977), volume 668 of Lecture Notes in Math., pages 231–244. Springer, Berlin, 1978.
- [36] B. Weiss, Sofic groups and dynamical systems, (Ergodic theory and harmonic analysis, Mumbai, 1999) *Sankhya Ser. A.* **62** (2000), 350–359.

DIPARTIMENTO DI INGEGNERIA, UNIVERSITÀ DEL SANNIO, 82100 BENEVENTO, ITALY
Email address: `tullio.cs@sbai.uniroma1.it`

UNIVERSITÉ DE STRASBOURG, CNRS, IRMA UMR 7501, F-67000 STRASBOURG, FRANCE
Email address: `michel.coornaert@math.unistra.fr`

DEPARTMENT OF MATHEMATICS, SUNY AT BUFFALO, BUFFALO, NY 14260-2900, USA
Email address: `hfli@math.buffalo.edu`