


This article was downloaded by:[University at Buffalo (SUNY)]
On: 19 October 2007
Access Details: [subscription number 769428387]
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number: 1072954
Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Communications in Algebra

Publication details, including instructions for authors and subscription information:
<http://www.informaworld.com/smpp/title~content=t713597239>

Symmetric Group Modules with Specht and Dual Specht Filtrations

David J. Hemmer^a

^a Department of Mathematics, University of Buffalo, SUNY, Buffalo, New York, USA

Online Publication Date: 01 November 2007

To cite this Article: Hemmer, David J. (2007) 'Symmetric Group Modules with Specht and Dual Specht Filtrations', Communications in Algebra, 35:11, 3292 - 3306

To link to this article: DOI: 10.1080/00927870701410611

URL: <http://dx.doi.org/10.1080/00927870701410611>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article maybe used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

SYMMETRIC GROUP MODULES WITH SPECHT AND DUAL SPECHT FILTRATIONS

David J. Hemmer

Department of Mathematics, University of Buffalo, SUNY,
Buffalo, New York, USA

The author and Nakano recently proved that multiplicities in a Specht filtration of a symmetric group module are well-defined precisely when the characteristic is at least five. This result suggested the possibility of a symmetric group theory analogous to that of good filtrations and tilting modules for $GL_n(k)$. This article is an initial attempt at such a theory. We obtain two sufficient conditions that ensure a module has a Specht filtration, and a formula for the filtration multiplicities. We then study the categories of modules that satisfy the conditions, in the process obtaining a new result on Specht module cohomology.

Next we consider symmetric group modules that have both Specht and dual Specht filtrations. Unlike tilting modules for $GL_n(k)$, these modules need not be self-dual, and there is no nice tensor product theorem. We prove a correspondence between indecomposable self-dual modules with Specht filtrations and a collection of $GL_n(k)$ -modules which behave like tilting modules under the tilting functor. We give some evidence that indecomposable self-dual symmetric group modules with Specht filtrations may be indecomposable self dual trivial source modules.

Key Words: Cohomology; Specht filtration; Symmetric group; Tilting module.

2000 Mathematics Subject Classification: Primary 20C30; Secondary 20G05.

1. INTRODUCTION

Let k be an algebraically closed field and let G be a reductive algebraic group over k . A rational G -module is said to have a good filtration if it has a filtration with successive quotients isomorphic to induced modules $\nabla(\lambda)$. There is a simple cohomological criterion for having a good filtration, and a formula for the multiplicities, which are independent of the choice of filtration. The indecomposable modules which have both a good and a Weyl filtration are called *tilting modules*, and are labelled by dominant weights.

Received April 26, 2006; Revised August 7, 2006. Communicated by D. K. Nakano.

Address correspondence to David J. Hemmer, Department of Mathematics, University of Buffalo, SUNY, Buffalo, NY 14260, USA; Fax: (419) 530-4720; E-mail: dhemmer@math.buffalo.edu

Until recently it was thought no such theory could exist for Specht and dual Specht filtrations of symmetric group modules, since well-known examples in characteristic two and three demonstrated that filtration multiplicities are not even well-defined. Nevertheless, in Hemmer and Nakano (2004) it was shown that the multiplicities are well-defined as long as $\text{char } k > 3$. Necessary and sufficient conditions for determining if a module has a Specht and/or dual Specht filtration were obtained. However the conditions are not in terms of symmetric group cohomology, but rather are stated in terms of GL_n cohomology and the adjoint Schur functor.

This article is a first attempt at a theory of Specht filtrations. We give two different sufficient conditions for a $k\Sigma_d$ module to have a Specht (or dual Specht) filtration. Although the conditions are not necessary, they have the advantage of being stated entirely in terms of the symmetric group theory. For modules satisfying the conditions, we obtain a formula for the filtration multiplicities which generalizes a known formula for Young modules. The collection of modules satisfying the condition gives an interesting subcategory of $\text{mod-}k\Sigma_d$, which we study.

We then consider modules which have both Specht and dual-Specht filtrations. We demonstrate that they are not as well-behaved as tilting modules. Unlike tilting modules, they need not be self-dual. The tensor product of two modules which have both Specht and dual Specht filtrations may have neither! We believe however that a classification of the indecomposable *self-dual* modules with both filtrations is possible. We show they are in correspondence with $GL_n(k)$ modules satisfying a certain natural property under the tilting functor. We give some evidence, inspired by recent work of Paget and Wildon, to suggest these modules may be self-dual trivial-source modules.

2. NOTATION AND PRELIMINARIES

We assume $\text{char } k > 3$ throughout, and emphasize that many of the results do not otherwise hold. Let $G = GL_n(k)$ and $V \cong k^n$ be the natural G -module. Let

$$S := S(n, d) \cong \text{End}_{k\Sigma_d}(V^{\otimes d})$$

be the Schur algebra. Modules for S correspond to polynomial G modules of homogeneous degree d . Our basic references for representation theory of $k\Sigma_d$ and of $S(n, d)$ are James (1978) and Martin (1993b).

Write $\lambda \vdash d$ for $\lambda = (\lambda_1, \dots, \lambda_r)$ a partition of d and let $\Lambda^+(d)$ be all partitions of d . Let $\Lambda^+(n, d)$ denote the partitions of d with at most n parts. Let $\Lambda(n, d)$ denote the compositions of d with at most n parts and let λ' denote the transpose of λ . Simple S -modules are indexed by $\Lambda^+(n, d)$ and denoted $L(\lambda)$. Let $\nabla(\lambda)$ and $\Delta(\lambda)$ denote the induced and Weyl modules, and $P(\lambda)$, $I(\lambda)$, and $T(\lambda)$ the projective, injective, and tilting modules for S . The tensor products of symmetric (resp., exterior) powers of V are denoted $S^\lambda(V)$ (resp., $\Lambda^\lambda(V)$). Descriptions of these modules can be found in Martin (1993b). Let $\mathcal{F}(\nabla)$ (resp., $\mathcal{F}(\Delta)$) be the set of S -modules having good (resp., Weyl) filtrations.

A partition λ is called p -restricted if $\lambda_i - \lambda_{i+1} < p$ for all i . It is p -regular if λ' is p -restricted. The simple $k\Sigma_d$ modules are indexed by p -restricted partitions and denoted D_λ . They can also be indexed by p -regular partitions and denoted D^μ .

The Specht, Young and permutation modules for $k\Sigma_d$ are indexed by $\Lambda^+(d)$ and denoted S^λ , Y^λ and M^λ . Let $S_\lambda = (S^\lambda)^*$. Recall that

$$\begin{aligned} S^\lambda \otimes \text{sgn} &\cong S_{\lambda'} \\ D^\lambda \otimes \text{sgn} &\cong D_{\lambda'}. \end{aligned} \quad (2.1)$$

2.1. Schur and Adjoint Schur Functors

For $n \geq d$ let $e \in S(n, d)$ denote the idempotent described in Green (1980, (6.1)). Then $eSe \cong k\Sigma_d$, and the Schur functor $\mathcal{F} : \text{mod-}S \rightarrow \text{mod-}k\Sigma_d$ is defined by $\mathcal{F}(U) := eU$. Let τ denote the usual contravariant duality on $\text{mod-}S$. Then \mathcal{F} is compatible with τ and the usual duality on $\text{mod-}k\Sigma_d$, i.e., $\mathcal{F}(U^\tau) \cong (\mathcal{F}(U))^*$.

The Schur functor is an exact, covariant functor with

$$\begin{aligned} \mathcal{F}(\nabla(\lambda)) &= S^\lambda & \mathcal{F}(\Delta(\lambda)) &= S_\lambda & \mathcal{F}(L(\lambda)) &= D_\lambda \text{ or } 0 \\ \mathcal{F}(P(\lambda)) &= Y^\lambda & \mathcal{F}(I(\lambda)) &= Y^\lambda & \mathcal{F}(T(\lambda)) &= Y^{\lambda'} \otimes \text{sgn}. \end{aligned} \quad (2.2)$$

The Schur functor admits a right adjoint functor $\mathcal{G} : \text{mod-}k\Sigma_d \rightarrow \text{mod-}S$ defined by

$$\begin{aligned} \mathcal{G}(N) &:= \text{Hom}_{k\Sigma_d}(V^{\otimes d}, N) \\ &\cong \text{Hom}_{eSe}(eS, N). \end{aligned}$$

The functor \mathcal{G} is a one-sided inverse to \mathcal{F} , i.e., $\mathcal{F}(\mathcal{G}(M)) \cong M$. The functor \mathcal{G} is only left exact, and so has higher right derived functors

$$R^i\mathcal{G}(N) = \text{Ext}_{k\Sigma_d}^i(V^{\otimes d}, N).$$

We now collect some known results about \mathcal{G} and $R^1\mathcal{G}$. In Doty et al. (2004) a Grothendieck spectral sequence is constructed to relate the cohomology of S and $k\Sigma_d$ using \mathcal{F} and \mathcal{G} . We will only use the related five-term exact sequence which begins

$$0 \rightarrow \text{Ext}_S^1(U, \mathcal{G}(N)) \rightarrow \text{Ext}_{k\Sigma_d}^1(\mathcal{F}(U), N) \rightarrow \text{Hom}_S(U, R^1\mathcal{G}(N)) \rightarrow \dots \quad (2.3)$$

Recall that we are assuming $p > 3$ throughout, indeed most of the results below fail for $p \leq 3$.

Proposition 2.1.

- (i) (Kleshchev and Nakano, 2001, 3.2). $\mathcal{G}(S_\lambda) \cong \Delta(\lambda)$.
- (ii) (Cline et al., 1996, 5.2.4). $\mathcal{G}(Y^\lambda \otimes \text{sgn}) \cong T(\lambda')$, $\mathcal{G}(M^\lambda \otimes \text{sgn}) \cong \Lambda^\lambda(V)$.
- (iii) (Hemmer and Nakano, 2004, 3.8.2). $\mathcal{G}(Y^\lambda) \cong P(\lambda)$, $\mathcal{G}(M^\lambda) \cong S^\lambda(V)$.
- (iv) (Hemmer and Nakano, 2004, 3.5.1). $M \in \text{mod-}k\Sigma_d$ has a dual Specht filtration if and only if $\mathcal{G}(M)$ has a Weyl filtration.
- (v) (Doty et al., 2004, 2.1.2). $\text{Hom}_S(L(\lambda), \mathcal{G}(M)) = 0$ unless λ is p -restricted. In particular if $L(\mu)$ lies in $\text{soc}(\Delta(\lambda))$, then μ must be p -restricted.

We will also need some information about $R^1\mathcal{G}$.

Proposition 2.2.

- (i) (Kleshchev and Nakano, 2001, 6.4). $R^1\mathcal{G}(S_\lambda) = 0$.
- (ii) If M has dual Specht filtration, then $\text{Ext}_S^1(U, \mathcal{G}(M)) \cong \text{Ext}_{k\Sigma_d}^1(\mathcal{F}(U), M)$.

Proof. Notice that (ii) follows from (i) and (2.3). □

3. A FILTRATION CRITERION AND MULTIPLICITY FORMULA

There is a well-known necessary and sufficient condition for an S module to have a good filtration and a formula for the filtration multiplicities.

Proposition 3.1 (Donkin, 1998, Proposition A2.2). *Let $V \in \text{mod-}S(n, d)$. Then:*

- (i) $V \in \mathcal{F}(\Delta)$ if and only if $\text{Ext}_S^1(V, \nabla(\lambda)) = 0$ for all $\lambda \in \Lambda^+(n, d)$. If so, then the multiplicity of $\Delta(\lambda)$ is independent of the choice of Weyl filtration and given by

$$[V : \Delta(\lambda)] = \dim_k \text{Hom}_S(V, \nabla(\lambda));$$

- (ii) $V \in \mathcal{F}(\nabla)$ if and only if $\text{Ext}_S^1(\Delta(\lambda), V) = 0$ for all $\lambda \in \Lambda^+(n, d)$. If so, then the multiplicity of $\nabla(\lambda)$ is independent of the choice of good filtration and given by

$$[V : \nabla(\lambda)] = \dim_k \text{Hom}_S(\Delta(\lambda), V).$$

Propositions 2.1(iv) and 3.1 immediately imply that a $k\Sigma_d$ module M has a dual Specht filtration if and only if $\text{Ext}_S^1(\mathcal{G}(M), \nabla(\lambda)) = 0$ for all $\lambda \in \Lambda^+(d)$. It seems unsatisfying to have a condition stated in terms of the S -cohomology of $\mathcal{G}(M)$ instead of the Σ_d -cohomology of M . The corresponding multiplicity formula is also in terms of S , namely,

$$[M : S_\mu] = \dim_k \text{Hom}_S(\mathcal{G}(M), \nabla(\mu)).$$

Given a $k\Sigma_d$ module with a Specht filtration, it is natural to ask if the filtration multiplicities are given by the dimension of some space of $k\Sigma_d$ -homomorphisms. There is one situation where this is known to be the case, namely, for Young modules in characteristic $p > 3$. It is well-known that Y^λ is self-dual with both a Specht and dual Specht filtration. The filtration multiplicities are well-defined in characteristic $p > 3$ by the main result of Hemmer and Nakano (2004), and they are given by the following proposition.

Proposition 3.2. *Let $[Y^\lambda : S_\mu]$ denote the multiplicity of S_μ in a dual Specht filtration of Y^λ . Then*

$$\begin{aligned} [Y^\lambda : S_\mu] &= \dim_k \text{Hom}_{k\Sigma_d}(Y^\lambda, S_\mu) \\ &= \dim_k \text{Hom}_{k\Sigma_d}(S^\mu, Y^\lambda). \end{aligned} \tag{3.1}$$

Proof. We show both sides of (3.1) are given by the decomposition number $[\Delta(\mu) : L(\lambda)]$. First observe that

$$\begin{aligned} [\Delta(\mu) : L(\lambda)] &= \dim_k \text{Hom}_S(P(\lambda), \Delta(\mu)) \\ &= \dim_k \text{Hom}_S(P(\lambda), \mathcal{G}(S_\mu)) \\ &= \dim_k \text{Hom}_{k\Sigma_d}(Y^\lambda, S_\mu) \quad \text{by adjointness.} \end{aligned}$$

But the decomposition number is also a filtration multiplicity in a tilting module:

$$\begin{aligned} [\Delta(\mu) : L(\lambda)] &= [T(\lambda') : \nabla(\mu')] \quad \text{by Donkin (1993, Lemma 3.1)} \\ &= [Y^\lambda \otimes \text{sgn} : S^\mu] \quad \text{since } \mathcal{F} \text{ is exact} \\ &= [Y^\lambda : S_\mu] \quad \text{by (2.1).} \quad \square \end{aligned}$$

More generally we can ask the following question.

Problem 3.3. Can one classify which $k\Sigma_d$ modules with Specht or dual Specht filtrations have multiplicities given by a formula like that in Proposition 3.2?

In Section 3.1 we give a large class of modules for which Problem 3.3 has an affirmative answer.

3.1. A Symmetric Group Filtration Criterion

We now prove that the symmetric group conditions corresponding to Proposition 3.1 under the Schur functor are sufficient to guarantee existence of filtrations. We obtain a pair of conditions which guarantee a Specht filtration and a pair which guarantee a dual Specht filtration. These are the first known conditions stated in terms of symmetric group cohomology that guarantee modules have certain filtrations. We also obtain a formula for the filtration multiplicities which generalizes Proposition 3.2. Notice that (i) and (iv) below correspond to Proposition 3.1 under \mathcal{F} .

Theorem 3.4. *Let $M \in \text{mod-}\Sigma_d$.*

- (i) *If $\text{Ext}_{k\Sigma_d}^1(M, S^\lambda) = 0 \ \forall \lambda \in \Lambda^+(d)$, then M has a dual Specht filtration. The multiplicity of S_μ in any such filtration is given by $\dim_k \text{Hom}_{k\Sigma_d}(M, S^\mu)$*
- (ii) *If $\text{Ext}_{k\Sigma_d}^1(S^\lambda, M) = 0 \ \forall \lambda \in \Lambda^+(d)$, then M has a dual Specht filtration. The multiplicity of S_μ in any such filtration is given by $\dim_k \text{Hom}_{k\Sigma_d}(S^\mu, M)$*
- (iii) *If $\text{Ext}_{k\Sigma_d}^1(M, S_\lambda) = 0 \ \forall \lambda \in \Lambda^+(d)$, then M has a Specht filtration. The multiplicity of S^μ in any such filtration is given by $\dim_k \text{Hom}_{k\Sigma_d}(M, S_\mu)$*
- (iv) *If $\text{Ext}_{k\Sigma_d}^1(S_\lambda, M) = 0 \ \forall \lambda \in \Lambda^+(d)$, then M has a Specht filtration. The multiplicity of S^μ in any such filtration is given by $\dim_k \text{Hom}_{k\Sigma_d}(S_\mu, M)$.*

Proof. We first prove (iv), so assume $\text{Ext}_{k\Sigma_d}^1(S_\lambda, M) = 0 \ \forall \lambda \in \Lambda^+(d)$. Then by (2.3) we get that $\text{Ext}_S^1(\Delta(\lambda), \mathcal{G}(M)) = 0 \ \forall \lambda \in \Lambda(d)$. By Proposition 3.1, $\mathcal{G}(M)$ has a good filtration, and the multiplicity of $\nabla(\mu)$ in $\mathcal{G}(M)$ is given by $\text{Hom}(\Delta(\mu), \mathcal{G}(M))$.

So $M = \mathcal{F}(\mathcal{G}(M))$ has a Specht filtration and the multiplicity formula follows from the adjointness of \mathcal{F} and \mathcal{G} . Now (i) follows immediately. To obtain (ii) and (iii) use (2.2) and Proposition 3.1. \square

We remark that the two criteria in each pair detect different modules. For instance a Young module Y^λ has both a Specht and dual Specht filtration. It is known that $\text{Ext}_{k\Sigma_d}^1(S^\lambda, Y^\mu)$ is always zero but $\text{Ext}_{k\Sigma_d}^1(Y^\mu, S^\lambda)$ may be nonzero (Hemmer and Nakano, 2004, 4.1), so the dual Specht filtration of Y^μ is detected by Theorem 3.4(ii) but not (i). Note that the multiplicity formula in Theorem 3.4(ii) is a significant generalization of Proposition 3.2.

The criteria in Theorem 3.4 are sufficient but not necessary; it is possible for a module to have a Specht filtration but not satisfy either Theorem 3.4(iii) or (iv). The structure of the Young and Specht modules for Σ_{2p} is well understood (see for example Martin, 1993a). From this one can easily calculate the following.

Example 3.5. Let $p = 5$ and $d = 10$. Then

$$\text{Ext}_{\Sigma_{10}}^1(S^{531^2}, S_{51^5}) \cong k \cong \text{Ext}_{\Sigma_{10}}^1(S_{55}, S^{531^2})$$

so the Specht filtration of S^{531^2} is not detected by either Theorem 3.4(iii) or (iv).

3.2. A Covariantly Finite Subcategory of $\text{mod-}k\Sigma_d$

The categories of $k\Sigma_d$ modules which satisfy the various conditions in Theorem 3.4 should be interesting to study. Let $\Theta = \{S^\lambda \mid \lambda \in \Lambda^+(d)\}$ and let $\mathcal{F}(\Theta)$ denote the full subcategory of $\text{mod } k\Sigma_d$ having Specht filtrations. $\mathcal{F}(\Theta)$ is obviously closed under extensions and it follows from Hemmer and Nakano (2004, 3.6.2) that it is closed under direct summands.

Lemma 3.6 (Hemmer and Nakano, 2004, 4.2.1). *Let $p > 3$ and suppose $\mu \not\prec \lambda$. Then $\text{Ext}_{k\Sigma_d}^1(S^\mu, S^\lambda) = 0$.*

Lemma 3.6 together with the main result of Ringel (1991) imply that $\mathcal{F}(\Theta)$ is functorially finite and has almost split sequences.

Observe that duality interchanges modules satisfying condition (i) with condition (iv) and (ii) with (iii) in Theorem 3.4. Tensoring with sgn interchanges those satisfying (i) with (iii) and (ii) with (iv). Thus if we want to study the module categories which arise from Theorem 3.4, we may without loss of generality consider Theorem 3.4(ii) and study the following category:

$$\mathcal{Y}(\Theta) := \{M \in \text{mod-}k\Sigma_d \mid \text{Ext}_{k\Sigma_d}^1(S^\lambda, M) = 0 \ \forall \lambda \in \Lambda^+(d)\}.$$

Notice that Theorem 3.4(ii) guarantees all modules in $\mathcal{Y}(\Theta)$ have dual Specht filtrations. Categories such as $\mathcal{Y}(\Theta)$ are studied in Ringel (1991) where, for example, it is shown that they are covariantly finite, i.e., every $k\Sigma_d$ -module has a left $\mathcal{Y}(\Theta)$ -approximation. These approximations are constructed explicitly in Ringel (1991, Sect. 2). For another example see Hemmer and Nakano (2004, Sect. 4), where the Young module Y^λ is constructed. The construction there actually builds Y^λ as the

left $\mathcal{Y}(\Theta)$ -approximation of S^λ , and can be applied for any $M \in \text{mod-}k\Sigma_d$ in a similar fashion.

By definition the Ext-injectives in the category $\mathcal{F}(\Theta)$ are modules $\mathcal{F}(\Theta) \cap \mathcal{Y}(\Theta)$. We next show the indecomposable objects in this set are exactly the Young modules.

Proposition 3.7. *The collection of indecomposable modules in $\mathcal{F}(\Theta) \cap \mathcal{Y}(\Theta)$ is exactly $\{Y^\lambda \mid \lambda \in \Lambda^+(d)\}$.*

Proof. Let $M \in \mathcal{F}(\Theta) \cap \mathcal{Y}(\Theta)$ be indecomposable, so M has a Specht filtration and $\forall \lambda \in \Lambda^+(d)$:

$$\begin{aligned} 0 &= \text{Ext}_{k\Sigma_d}^1(S^\lambda, M) \\ &\cong \text{Ext}_{k\Sigma_d}^1(S_{\lambda'}, M \otimes \text{sgn}) \\ &\cong \text{Ext}_S^1(\Delta(\lambda'), \mathcal{G}(M \otimes \text{sgn})) \quad \text{by Proposition 2.2(ii)}. \end{aligned}$$

Thus $\mathcal{G}(M \otimes \text{sgn})$ is indecomposable and has a good filtration by Proposition 3.1. But $M \otimes \text{sgn}$ has a dual Specht filtration so $\mathcal{G}(M \otimes \text{sgn})$ has a Weyl filtration by Proposition 2.1. So $\mathcal{G}(M \otimes \text{sgn})$ has both a good and Weyl filtration and hence is isomorphic to a tilting module. So $M \otimes \text{sgn} \cong \mathcal{F}(T(\lambda))$ for some λ , so $M \cong Y^\lambda$ by (2.2). □

A similar argument shows the indecomposable Ext-projectives in the category $\mathcal{F}(\Theta)$ are the twisted Young modules $\{Y^\lambda \otimes \text{sgn}\}$. We remark that the set $\{(S^\lambda, Y^\lambda)\}$ has recently been shown to be a *stratifying system*, see Erdmann (2005). This result together with Theorem 2.4 of Marcos et al. (2004) could also be used to prove Proposition 3.7.

It would be nice to understand exactly which modules with dual Specht filtrations are in $\mathcal{Y}(\Theta)$. Of course the simplest module with a dual Specht filtration is just a dual Specht module, and in this case we can give a necessary and sufficient condition for $S_\mu \in \mathcal{Y}(\Theta)$.

Proposition 3.8. *$S_\mu \in \mathcal{Y}(\Theta)$ if and only if:*

- (a) $\mathcal{G}(S^\mu) \cong \nabla(\mu')$ and
- (b) $R^1\mathcal{G}(S^\mu) = 0$.

Proof. Suppose $S_\mu \in \mathcal{Y}(\Theta)$, so $0 = \text{Ext}_{k\Sigma_d}^1(S^\tau, S_\mu) = \text{Ext}_{k\Sigma_d}^1(S_\tau, S^\mu)$ for all $\tau \in \Lambda^+(d)$. From (2.3) we get that $\text{Ext}^1(\Delta(\tau), \mathcal{G}(S^\mu)) = 0 \forall \tau$. Thus $\mathcal{G}(S^\mu)$ has a good filtration and maps to S^μ under \mathcal{F} . Since the multiplicities in a Specht filtration are well-defined we must have $\mathcal{G}(S^\mu) \cong \nabla(\mu')$. Furthermore, $V^{\otimes d}$ has a dual Specht filtration, so $\text{Ext}_{k\Sigma_d}^1(V^{\otimes d}, S^\mu) = R^1\mathcal{G}(S^\mu) = 0$.

Conversely, assume $\mathcal{G}(S^\mu) \cong \nabla(\mu')$ and $R^1\mathcal{G}(S^\mu) = 0$. Plugging into (2.3) immediately implies $\text{Ext}_{k\Sigma_d}^1(S_\tau, S^\mu) = 0 \forall \tau \in \Lambda^+(d)$, so $S_\mu \in \mathcal{Y}(\Theta)$. □

It seems a difficult question to determine for which μ will the conditions of Proposition 3.8 hold.

Since $\text{soc}(\nabla(\mu')) \cong L(\mu')$, Propositions 2.1(iv), (v) imply the conditions of Proposition 3.8 hold only if μ' is p -restricted, i.e., if μ is p -regular. A similar argument shows that S_μ is detected by Theorem 3.4(i) only if μ is p -restricted. Thus we have obtained some new information about extensions between Specht and dual Specht modules.

Corollary 3.9.

- (i) If μ is not p -regular, then $\text{Ext}_{k\Sigma_d}^1(S^\lambda, S_\mu) \neq 0$ for some $\lambda \in \Lambda^+(d)$.
- (ii) If μ is not p -restricted, then $\text{Ext}_{k\Sigma_d}^1(S_\mu, S^\lambda) \neq 0$ for some $\lambda \in \Lambda^+(d)$.
- (iii) If μ is neither p -regular nor p -restricted, then the obvious dual Specht filtration of S_μ is not detected by Theorem 3.4(i) or (ii).

Setting $\mu = (d)$ in Corollary 3.9(ii) gives new information about Specht module cohomology.

Corollary 3.10. *Let $d \geq p$. Then there exists some $\lambda \in \Lambda^+(d)$ with $H^1(\Sigma_d, S^\lambda) \neq 0$.*

The previous corollary is in stark contrast to the situation for dual Specht modules where in char $p > 3$, it is known (Burichenko et al., 1996) that $H^1(\Sigma_d, S_\lambda) = 0$ for all λ .

4. A REVEALING EXAMPLE

The modules in $\mathcal{F}(\Delta) \cap \mathcal{F}(\nabla)$ are called *tilting modules*. For each dominant weight λ there is a unique indecomposable tilting module $T(\lambda)$ with highest weight λ . These modules are self-dual and an arbitrary tilting module is a direct sum of them. Since $\mathcal{F}(\nabla)$ and thus $\mathcal{F}(\Delta)$ are closed under tensor products (Jantzen, 2003, II.4.21), then so is the collection of tilting modules (as G modules not S modules). Tilting modules also have nice cohomological properties. For example $\text{Ext}_S^i(T(\lambda), T(\mu)) = 0 \forall i > 0$.

In this section we give an example which shows how these properties fail for symmetric group modules with both Specht and dual Specht filtrations. Later, we conjecture that a nice theory may be salvaged if one assumes additionally that the modules are self-dual.

Let $p = d = 5$. The module category for $k\Sigma_p$ in characteristic p is completely understood, indeed has only finitely many indecomposable modules.

Example 4.1. Let $U \cong \Omega^2(D_{21^3})$ where Ω is the Heller translate. Then U has the following structure:



The Specht modules in the principal block have the following Loewy structures:

$$S^5 \cong D_{41}, \quad S^{41} \cong \begin{array}{c} D_{31^2} \\ \hline D_{41} \end{array}, \quad S^{31^2} \cong \begin{array}{c} D_{21^3} \\ \hline D_{31^2} \end{array}, \quad S^{21^3} \cong \begin{array}{c} D_{1^5} \\ \hline D_{21^3} \end{array}, \quad S^{1^5} \cong D_{1^5}, \quad (4.1)$$

so U has a Specht filtration with subquotients S^{1^5} , S^{31^2} , and S^5 and a dual Specht filtration with subquotients S_{41} and S_{21^3} .

Proposition 4.2. *Let $U \cong \Omega^2(D_{21^3})$. Then:*

- (i) $U \otimes U$ has neither a Specht nor dual Specht filtration;
- (ii) $\text{Ext}_{\Sigma_5}^1(U, U^*) \cong k$;
- (iii) U does not lift to characteristic zero;
- (iv) U has vertex $P \in \text{Syl}_5(\Sigma_5)$. It has source isomorphic to the unique indecomposable kP module of Loewy length three, i.e., $\text{rad}^2(kP)$.

Proof. We have

$$\begin{aligned} U \otimes U &\cong \Omega^2(D_{21^3}) \otimes \Omega^2(D_{21^3}) \\ &\cong \Omega^4(D_{21^3} \otimes D_{21^3}) \oplus P \\ &\cong \Omega^4(D_{21^3} \oplus D_{41} \oplus D_{32}) \oplus P \\ &\cong D_{31^2} \oplus D_{1^5} \oplus P' \end{aligned}$$

where P, P' are projective modules. We used the fact that D_{32} is projective and that the projective resolutions of the simple modules are easy to write down. Also the structure $D_{21^3} \otimes D_{21^3}$ can be easily computed by hand or computer, as D_{21^3} is only three dimensional, the details are left to the reader. Since projectives have both filtrations and $D_{31^2} \oplus D_{1^5}$ has neither, we conclude that $U \otimes U$ has neither.

To prove (ii) observe that $U^* \cong \Omega^2(D_{31^2})$ so $\text{Ext}^1(U, U^*) \cong \text{Ext}^1(D_{21^3}, D_{31^2}) \cong k$. Next notice that if U could be lifted to characteristic zero then so could D_{21^3} , which from (4.1) is not the case. The last part can also be computed by hand. In particular,

$$\text{Ind}_P^{\Sigma_5} \text{rad}^2(kP) \cong U \oplus U^* \oplus D_{21^3} \oplus D_{31^2}. \quad \square$$

This example easily generalizes to Σ_p for any $p \geq 5$. None of the similar “pathological” examples which we have constructed have been indecomposable and self-dual. Thus in the next section we consider these modules.

5. “TILTING” MODULES FOR SYMMETRIC GROUPS?

Despite the example in the previous section, we believe that assuming an indecomposable module with both Specht and dual Specht filtrations is also self-dual might allow a complete classification, and we further believe that the geometric tools available in the algebraic group theory could be useful. To further this goal, we prove that these modules correspond bijectively with a nice class of G -modules. Roughly speaking these modules behave like tilting modules under the tilting functor in a way we make precise below. If our conjecture in the next section is correct, then these modules will be indecomposable self-dual trivial source modules.

5.1. Ringel Duals and Tilting Functors

Let $T = \bigoplus_{\alpha \in \Lambda(n,d)} (\Lambda^\alpha(V))$. Then $T \cong T^\tau$ is a full tilting module and the Ringel dual of $S(n, d)$ is defined as

$$S'(n, d) \cong \widetilde{\text{End}}_{S(n,d)}(T).$$

The tilting functor $\mathcal{F} : \text{mod-}S(n, d) \rightarrow \text{mod-}S'(n, d)$ is given by

$$\mathcal{F}(U) = \text{Hom}_S(T, U).$$

This setup is more thoroughly described in Donkin (1993). When $n \geq d$, then Donkin (1993, Sect. 3) proved that

$$S(n, d) \cong \text{End}_{k\Sigma_d} \left(\bigoplus_{\alpha \in \Lambda(n,d)} M^\alpha \right), \quad S'(n, d) \cong \text{End}_{k\Sigma_d} \left(\bigoplus_{\alpha \in \Lambda(n,d)} M^\alpha \otimes \text{sgn} \right). \quad (5.1)$$

In particular $S(n, d) \cong S'(n, d)$. We let $\widetilde{\mathcal{F}} : \text{mod-}S(n, d) \rightarrow \text{mod-}S(n, d)$ denote the composition of \mathcal{F} with the functor $\text{mod-}S'(n, d) \rightarrow \text{mod-}S(n, d)$ arising from the isomorphism. This is the functor denoted $\widetilde{\mathcal{F}}$ in Donkin (1993). It is well-known that \mathcal{F} , and thus $\widetilde{\mathcal{F}}$, takes modules with good filtrations to modules with Weyl filtrations and interchanges projective and tilting modules.

Lemma 5.1 (Donkin, 1993, Section 3).

- (i) $\widetilde{\mathcal{F}}(\nabla(\lambda)) \cong \Delta(\lambda')$, $\widetilde{\mathcal{F}}(\mathcal{F}(\nabla)) \subseteq \mathcal{F}(\Delta)$.
- (ii) $\widetilde{\mathcal{F}}(T(\lambda)) \cong P(\lambda')$.
- (iii) $\widetilde{\mathcal{F}}(P(\lambda)) \cong T(\lambda')$.
- (iv) $\widetilde{\mathcal{F}}(I(\lambda)) \cong T(\lambda')$.

In order to relate $\widetilde{\mathcal{F}}$ to modules for $k\Sigma_d$ we need to express $\widetilde{\mathcal{F}}$ in terms of the \mathcal{F} and \mathcal{G} .

Proposition 5.2. *Let $U \in \text{mod-}S(n, d)$. Then $\widetilde{\mathcal{F}}(U) \cong \mathcal{G}(eU \otimes \text{sgn})$.*

Proof.

$$\begin{aligned} \mathcal{F}(U) &\cong \text{Hom}_{S(n,d)}(T, U) \\ &\cong \text{Hom}_{S(n,d)}(U^\tau, T) \\ &\cong \text{Hom}_{k\Sigma_d}((eU)^*, \bigoplus (M^\alpha \otimes \text{sgn})) \quad \text{since } \mathcal{G}(M^\alpha \otimes \text{sgn}) \cong \Lambda^\alpha(V) \\ &\cong \text{Hom}_{k\Sigma_d}(\bigoplus (M^\alpha \otimes \text{sgn}), eU). \end{aligned}$$

Now $\mathcal{F}(U)$ is a module for $S(n, d)'$ via its right action on $\bigoplus (M^\alpha \otimes \text{sgn})$. To determine $\widetilde{\mathcal{F}}(U)$ we need to compose with the isomorphism from (5.1), which

simply takes a map ϕ to $\phi \otimes \text{id}$. Thus we obtain

$$\begin{aligned}\tilde{\mathcal{F}}(U) &\cong \text{Hom}_{k\Sigma_d}(\oplus M^{\alpha}, eU \otimes \text{sgn}) \\ &\cong \text{Hom}_{k\Sigma_d}(V^{\otimes d}, eU \otimes \text{sgn}) \\ &\cong \mathcal{G}(eU \otimes \text{sgn})\end{aligned}$$

as desired. \square

This alternate description of $\tilde{\mathcal{F}}$ is all we need to prove.

Theorem 5.3. *Let $M \in \text{mod-}k\Sigma_d$ be indecomposable and let $U = \mathcal{G}(M)$.*

- (i) *M has both a Specht and dual Specht filtration if and only if both U and $\tilde{\mathcal{F}}(U)$ have Weyl filtrations.*
- (ii) *M is self dual with a Specht (and hence dual Specht) filtration if and only if U has a Weyl filtration and $\tilde{\mathcal{F}}(U) \cong \tilde{\mathcal{F}}(U^{\tau})$.*

Proof. Notice that $U \in \mathcal{F}(\Delta)$ implies $U^{\tau} \in \mathcal{F}(\nabla)$ which implies $\mathcal{F}(U^{\tau}) \in \mathcal{F}(\Delta)$ by Lemma 5.1. Thus the condition $\tilde{\mathcal{F}}(U) \cong \tilde{\mathcal{F}}(U^{\tau})$ from part (ii) implies (as expected) the condition from part (i) that $\tilde{\mathcal{F}}(U)$ have a Weyl filtration.

We use Proposition 5.2 repeatedly. First suppose U and $\tilde{\mathcal{F}}(U)$ both have Weyl filtrations. Then $M \cong eU$ has a dual Specht filtration as does $e\tilde{\mathcal{F}}(U) \cong M \otimes \text{sgn}$. Thus by (2.2), M has both filtrations. Conversely, if $M \in \text{mod-}k\Sigma_d$ is indecomposable with both filtrations then so is $M \otimes \text{sgn}$ and setting $U = \mathcal{G}(M)$ we immediately get that both U and $\tilde{\mathcal{F}}(U)$ have Weyl filtrations by Proposition 2.1(iv).

Now further suppose that $M \cong M^*$ is indecomposable with both filtrations and let $U = \mathcal{G}(M)$. Then $\tilde{\mathcal{F}}(U) \cong \mathcal{G}(M \otimes \text{sgn})$. Similarly, $\tilde{\mathcal{F}}(U^{\tau}) \cong \mathcal{G}(eU^{\tau} \otimes \text{sgn}) \cong \mathcal{G}(M^* \otimes \text{sgn})$ so $\tilde{\mathcal{F}}(U) \cong \tilde{\mathcal{F}}(U^{\tau})$. The same argument shows that $\tilde{\mathcal{F}}(U) \cong \tilde{\mathcal{F}}(U^{\tau})$ implies M is self-dual. \square

Remark 5.4. Observe from Lemma 5.1 that the tilting modules $U = T(\lambda)$ and the projective modules $U = P(\lambda)$ both have Weyl filtrations and satisfy $\tilde{\mathcal{F}}(U) \cong \tilde{\mathcal{F}}(U^{\tau})$, so the modules described in Theorem 5.3 are a simultaneous generalization of tilting modules and projective modules.

When $n < d$, then $S(n, d)$ and $S(n, d)'$ are usually not isomorphic and so the functor $\tilde{\mathcal{F}}$ does not exist, but the tilting functor \mathcal{F} does. So one can ask more generally the following question.

Problem 5.5. Which $S(n, d)$ modules U (or which $U \in \mathcal{F}(\Delta)$) have the property that $\mathcal{F}(U) \cong \mathcal{F}(U^{\tau})$?

In Hemmer (2006) we conjectured that indecomposable self-dual $k\Sigma_d$ -modules with Specht filtrations would be exactly signed Young modules. Equivalently, this conjecture says the modules in Theorem 5.3 would be the listing modules recently defined by Donkin. A counterexample has recently been found (Paget, 2007; Wildon, 2004).

Example 5.6 (Paget, 2007; Wildon, 2004). Let $H \cong C_2^p \rtimes \Sigma_p \leq \Sigma_{2p}$ be the normalizer of a fixed-point-free involution in Σ_{2p} and let $M \cong \text{Ind}_H^{\Sigma_{2p}} k$. Then each indecomposable summand of M is self-dual with a Specht filtration. One of these is not a signed Young module.

Specifically, Paget (2007) proved that the indecomposable summands of M have Specht filtrations, but Wildon (2004) had already verified that each summand of M is self-dual. It follows by general theory that one of the summands must have vertex a Sylow subgroup $P \leq H$. However P is not conjugate to the Sylow subgroup of any Young subgroup of Σ_{2p} , so Paget concluded this summand is not a signed Young module. We will discuss this situation further in Section 7.

6. IRREDUCIBLE SPECHT MODULES

Since the irreducible modules for $k\Sigma_d$ are self-dual, the irreducible Specht modules give an immediate source of modules with both Specht and dual Specht filtrations. For λ p -regular or p -restricted the criterion for S^λ to be irreducible has long been known. Only recently Fayers (2005) has verified a conjecture of James and Mathas which settles the problem completely. In this section we determine which of these modules are detected by the criteria from Section 3.

Lemma 6.1. *Suppose μ is p -regular and S^μ is irreducible. Then:*

- (i) $S^\mu \cong Y^\mu$;
- (ii) $\text{Ext}_{k\Sigma_d}(S^\lambda, S_\mu) = 0 \ \forall \lambda \in \Lambda^+(d)$;
- (iii) $\text{Ext}_{k\Sigma_d}(S_\mu, S_\lambda) = 0 \ \forall \lambda \in \Lambda^+(d)$.

Proof. Since μ is p -regular, then $S^\mu \cong D^\mu$. Also $[Y^\lambda : D^\lambda] = 1$. But Y^λ is self-dual and indecomposable so $Y^\lambda \cong D^\lambda \cong S^\lambda$. Parts (ii) and (iii) follow from the fact that $\text{Ext}^1(S^\lambda, Y^\mu) = 0$ for $p > 3$ and S^μ is self-dual (Hemmer and Nakano, 2004, 4.1.1). □

Notice parts (ii) and (iii) are saying that both the Specht and dual Specht filtrations of S^λ are detected by Theorem 3.4. Similarly we have the following lemma.

Lemma 6.2. *Suppose μ is p -restricted and S^μ is irreducible. Then:*

- (i) $S^\mu \cong Y^{\mu'} \otimes \text{sgn}$;
- (ii) $\text{Ext}_{k\Sigma_d}(S_\lambda, S_\mu) = 0 \ \forall \lambda \in \Lambda^+(d)$;
- (iii) $\text{Ext}_{k\Sigma_d}(S_\mu, S^\lambda) = 0 \ \forall \lambda \in \Lambda^+(d)$.

Proof. This follows from Lemma 6.1 immediately since $S^\mu \otimes \text{sgn} \cong S_{\mu'} \cong S^{\mu'}$ and μ' is p -regular. □

In stark contrast to the previous two lemmas, the next result shows that irreducible Specht modules which are neither p -restricted nor p -regular are *never* detected by Theorem 3.4.

Proposition 6.3. *Suppose μ is neither p -regular nor p -restricted and suppose S^μ is irreducible. Then S^μ does not satisfy any of the criteria in Theorem 3.4.*

Proof. Suppose S^μ satisfies the criterion in Theorem 3.4(ii), so $\text{Ext}_{k\Sigma_d}^1(S^\lambda, S^\mu) = 0 \forall \lambda$. Then $S^\mu \cong Y^\mu$ by Proposition 3.7. So $\mathcal{G}(S^\mu) \cong \Delta(\mu) \cong P(\mu)$. Thus $\Delta(\mu) \cong L(\mu)$ is an irreducible projective module and μ must be p -restricted, contradicting our assumption. Thus S^μ cannot satisfy criterion (ii). Similar arguments handle the other three criteria. \square

7. TRIVIAL SOURCE MODULES

Recall that the Young modules Y^λ are precisely the indecomposable summands of the permutation modules $M^\lambda \cong \text{Ind}_{\Sigma_\lambda}^{\Sigma_d} k$. For $\lambda \vdash a$ and $\mu \vdash b$ with $a + b = d$ define the *signed permutation module*

$$M^{(\lambda|\mu)} \cong \text{Ind}_{\Sigma_\lambda \times \Sigma_\mu}^{\Sigma_d} k \boxtimes \text{sgn}.$$

Indecomposable summands of signed permutation modules are called *signed Young modules*. Notice the class of signed Young modules includes all ordinary Young modules Y^λ and twisted Young modules $Y^\lambda \otimes \text{sgn}$ but, when $d \geq 2p$, includes other modules as well. Signed Young modules give a large class of self-dual modules with both Specht and dual Specht filtrations. We once believed this was exactly the class of indecomposable self-dual modules with Specht filtrations. However, Example 5.6 shows there are more.

The signed Young modules have trivial source and vertex isomorphic to a Sylow subgroup of some Young subgroup of Σ_d . However, in general there are many conjugacy classes of p -subgroups of Σ_d which are not of this form, and many more trivial source modules, for example the module M in Example 5.6. On the other hand the module U from Proposition 4.2 has both Specht and dual Specht filtrations, and is not a trivial-source module. But it is not self-dual either! Undaunted, we dare to hazard another conjecture.

Conjecture 7.1. *Indecomposable, self-dual $k\Sigma_d$ modules with Specht filtrations are trivial source modules.*

If this conjecture were true it would have the nice consequence that there are only finitely many indecomposable self-dual $k\Sigma_d$ modules with Specht filtrations.

As evidence for this we showed the following theorem.

Theorem 7.2 (Hemmer, 2006, Theorem 4.2). *Suppose S^μ be irreducible. Then S^μ is a signed Young module, and hence has trivial source.*

8. FINAL REMARKS AND PROBLEMS

Even though many of the results and the machinery of this article depend on the characteristic being at least five, it is possible that Conjecture 7.1 holds in characteristics two and three. For example in characteristic two, $d = 4$, there are

two irreducible Σ_4 -modules and both of them are isomorphic to Specht modules and both have trivial source. Thus *every* Σ_4 module has both a Specht and dual Specht filtration. For Conjecture 7.1 to hold here, it would require $k\Sigma_4$ to have only finitely many indecomposable self-dual modules, even though the algebra has wild type!

We close with a list of problems for further study.

Problem 8.1. Suppose M and N are indecomposable self-dual modules with Specht filtrations. Is $M \otimes N$ a direct sum of such modules?

We have been unable to answer this even for $M \otimes M$ where M is from Example 5.6.

Problem 8.2. Must indecomposable, trivial source modules have either Specht or dual Specht filtrations? More generally, must $k\Sigma_d$ modules which are p -modular reductions of some indecomposable \mathcal{O}_{Σ_d} -lattice have Specht or dual Specht filtrations?

For Σ_p this is known, the trivial source modules are exactly the Young and twisted Young modules, together with the Specht modules $S^{(p-a, 1^a)}$ and their duals $S_{(p-a, 1^a)}$ for a even (Erdmann, 2001, 4.3).

Problem 8.3. Suppose M and N are indecomposable self-dual modules with Specht filtrations and suppose $p > 3$. Is $\text{Ext}_{k\Sigma_d}^i(M, N) = 0$ for $1 \leq i \leq p - 3$?

This is known to be the case for signed Young modules (Hemmer, 2006, Lemma 2.2(iii)).

Notice that U in Example 4.1 does not lift to characteristic zero, by Proposition 4.2(iii). This is true of other indecomposable modules we have found which have both Specht and dual Specht filtrations but are not self dual. The signed Young modules do, however, lift to characteristic zero, as does any trivial source module.

Problem 8.4. Suppose M and N are indecomposable self-dual modules with Specht filtrations and suppose $p > 3$. Do M and N lift to characteristic zero?

If, for example, Problem 8.3 were answered affirmatively, it would in particular show that $\text{Ext}_{k\Sigma_d}^2(M, M) = 0$. This Ext^2 -vanishing would imply that M lifts to characteristic zero (Benson, 1991, 3.7.7). We remark that the module itself lifting to an integral representation is a weaker condition than the entire filtration lifting.

Notice that U from Example 4.1 does not self-extend, i.e., $\text{Ext}_{k\Sigma_d}^1(U, U) = 0$. We know of no examples of indecomposable modules with Specht and dual Specht filtrations that self-extend.

Problem 8.5. Suppose $U \in \text{mod-}k\Sigma_d$ is indecomposable with both Specht and dual Specht filtrations. Must $\text{Ext}_{k\Sigma_d}^1(U, U) = 0$?

ACKNOWLEDGMENTS

Author's research was supported in part by an NSA Young Investigator's Grant.

REFERENCES

- Benson, D. J. (1991). *Representations and Cohomology I: Basic Representation Theory of Finite Groups and Associative Algebras*. Cambridge Studies in Advanced Mathematics. Vol. 30. Cambridge: Cambridge University Press.
- Burichenko, V. P., Kleshchev, A. S., Martin, S. (1996). On cohomology of dual Specht modules. *J. Pure Appl. Algebra* 112(2):157–180.
- Cline, E., Parshall, B., Scott, L. (1996). Stratifying endomorphism algebras. *Mem. Amer. Math. Soc.* 124:591.
- Donkin, S. (1993). On tilting modules for algebraic groups. *Math. Z.* 212(1):39–60.
- Donkin, S. (1998). *The q -Schur Algebra*. London Mathematical Society Lecture Note Series. Vol. 253. Cambridge: Cambridge University Press.
- Doty, S. R., Erdmann, K., Nakano, D. K. (2004). *Extensions of Modules over Schur Algebras, Symmetric Groups, and Hecke Algebras*. Algebras and Representation Theory.
- Erdmann, K. (2001). Young modules for symmetric groups. *J. Austral. Math. Soc.* 71: 201–210.
- Erdmann, K. (2005). Stratifying systems, filtration multiplicities and symmetric groups. *J. Algebra Appl.* 4:551–555.
- Fayers, M. (2005). Irreducible Specht modules for Hecke algebras of type A. *Adv. Math.* 193:438–452.
- Green, J. A. (1980). *Polynomial Representations of GL_n* . Lecture Notes in Mathematics. Vol. 830. Berlin: Springer-Verlag.
- Hemmer, D. J. (2006). Irreducible Specht modules are signed Young modules. *J. Algebra* 305:433–441.
- Hemmer, D. J., Nakano, D. K. (2004). Specht filtrations for Hecke algebras of type A. *J. London Math. Soc.* 69.
- James, G. (1978). *The Representation Theory of the Symmetric Groups*. Lecture Notes in Math. Vol. 682. Springer-Verlag.
- Jantzen, J. C. (2003). *Representations of Algebraic Groups*. 2nd ed. Mathematical Surveys and Monographs. Vol. 107. American Mathematical Society.
- Kleshchev, A. S., Nakano, D. K. (2001). On comparing the cohomology of general linear and symmetric groups. *Pacific J. Math.* 201(2):339–355.
- Marcos, E. N., Mendoza, O., Sáenz, C. (2004). Stratifying systems via relative simple modules. *J. Algebra* 280.
- Martin, S. (1993a). Projective indecomposable modules for symmetric groups. I, *Quart. J. Math. Oxford Ser. (2)* 44(173):87–99.
- Martin, S. (1993b). *Schur Algebras and Representation Theory*. Cambridge Tracts in Mathematics. Vol. 112. Cambridge: Cambridge University Press.
- Paget, R. (2007). A family of modules with Specht and dual Specht filtrations. *J. Algebra* 312:880–890.
- Ringel, C. M. (1991). The category of modules with good filtrations over a quasi-hereditary algebra has almost split sequences. *Math. Z.* 208(2):209–223.
- Wildon, M. (2004). *Modular Representations of Symmetric Groups*. Ph.D. thesis, University College Oxford.