

# SMOOTH APPROXIMATION OF LIPSCHITZ PROJECTIONS

HANFENG LI

ABSTRACT. We show that any Lipschitz projection-valued function  $p$  on a connected closed Riemannian manifold can be approximated uniformly by smooth projection-valued functions  $q$  with Lipschitz constant close to that of  $p$ . This answers a question of Rieffel.

## 1. INTRODUCTION

The question of approximating continuous functions on Riemannian manifolds by smooth functions preserving geometric properties has a long history. In [1, 2, 3] Greene and Wu studied such questions for real-valued functions and geometric properties such as having Lipschitz constant bounded above by a fixed number, and used such results to obtain geometric applications [2].

In a previous version of [5] Rieffel asked the question whether, for any Lipschitz function  $p$  on a compact Riemannian manifold  $M$  with values in the projection set of a matrix algebra  $M_n(\mathbb{C})$  and any  $\varepsilon > 0$ , there is a smooth function  $q$  on  $M$  also with values in the projection set of  $M_n(\mathbb{C})$  with  $\|p - q\|_\infty < \varepsilon$  and  $L(q) < L(p) + \varepsilon$ . Here  $\|f\|_\infty$  denotes the supremum norm of  $f$  and

$$(1) \quad L(f) = \sup_{x, y \in M, x \neq y} \frac{\|f(x) - f(y)\|}{\rho(x, y)}$$

denotes the Lipschitz constant of  $f$ , for  $\rho$  denoting the geodesic distance on  $M$ . An affirmative answer to this question has direct application on obtaining lower bounds of the Lipschitz constants for projection-valued functions on  $M$  representing a fixed vector bundle on  $M$ .

In this note we answer Rieffel's question affirmatively. In fact, we shall deal more generally with functions with values in the projection set of any  $C^*$ -algebra. Following [5], by a *real  $C^*$ -subring* we mean a norm-closed  $*$ -subring of a  $C^*$ -algebra which is closed under multiplication by scalars in  $\mathbb{R}$ . For example,  $M_n(\mathbb{R})$  is a real  $C^*$ -subring

---

*Date:* October 14, 2009.

Partially supported by NSF Grant DMS-0701414.

contained in  $M_n(\mathbb{C})$ . For a compact manifold  $M$  and a real  $C^*$ -subring  $A$ , denote by  $C(M, A)$  the real  $C^*$ -subring of all continuous  $A$ -valued functions on  $M$ , and denote by  $C^\infty(M, A)$  the subalgebra of all smooth  $A$ -valued functions on  $M$ .

**Theorem 1.1.** *Let  $M$  be a connected closed Riemannian manifold. Let  $A$  be a real  $C^*$ -subring. For any projection  $p \in C(M, A)$  and any  $\varepsilon > 0$ , there exists a projection  $q \in C^\infty(M, A)$  with  $\|p - q\|_\infty < \varepsilon$  and  $L(q) < L(p) + \varepsilon$ .*

As a sample, Rieffel has obtained lower bound of the Lipschitz constants for projections representing certain complex line bundle on the two-torus  $\mathbb{T}^2$  in [5, Proposition 12.1], via obtaining such bound for smooth projections first and then extend the bound to nonsmooth projections using Theorem 1.1.

The proof of Theorem 1.1 has two ingredients. The first one is that part of Greene and Wu's results hold generally for functions valued in any Banach space. We recall these results in Section 2. The second one is a fine estimate of the Lipschitz seminorm of the projection in a  $C^*$ -algebra obtained in the usual way from an "almost-projection". We give this estimate and prove Theorem 1.1 in Section 3.

I am grateful to Wei Wu for comments.

## 2. SMOOTH APPROXIMATION OF CONTINUOUS FUNCTIONS

In [1] Greene and Wu introduced *the Riemannian convolution smoothing process* for approximating continuous real-valued functions on a Riemannian manifold by smoothing functions preserving geometric properties. They used this technique to obtain certain geometric application [2]. In fact, much of Greene and Wu's results work for functions valued in a Banach space. Let us recall their construction on [1, Page 646-647].

Fix a positive integer  $n$ . Let  $\kappa : \mathbb{R} \rightarrow \mathbb{R}$  be a non-negative smooth function which has support contained in the interval  $[-1, 1]$ , is constant in a neighborhood of 0, and satisfies  $\int_{v \in \mathbb{R}^n} \kappa(\|v\|) = 1$ , where  $\mathbb{R}^n$  is equipped with the standard Lebesgue measure. Let  $V$  be a real Banach space. Let  $M$  be a Riemannian manifold of dimension  $n$  without boundary, and let  $K$  be a compact subset of  $M$ . Then there is an  $\varepsilon_K > 0$  such that, for every  $x$  in some neighborhood of  $K$  and every  $v$  in the tangent space  $\text{TM}_x$  of  $M$  at  $x$  with  $\|v\| < \varepsilon_K$ , the exponential  $\exp_x v$  is defined. For any continuous function  $f : M \rightarrow V$  and any  $0 < \varepsilon < \varepsilon_K$  define a  $V$ -valued function  $f_\varepsilon$  on a neighborhood of  $K$  by

$$f_\varepsilon(x) = \frac{1}{\varepsilon^n} \int_{v \in \text{TM}_x} \kappa\left(\frac{\|v\|}{\varepsilon}\right) f(\exp_x v),$$

where  $TM_x$  is equipped with the standard Lebesgue measure via a linear isometry between  $TM_x$  and the Euclidean space  $\mathbb{R}^n$ .

In [1, Page 647] and [2, Lemma 8] Greene and Wu proved the following theorem for the case  $V = \mathbb{R}$ . Their proof works for any Banach space  $V$ .

**Theorem 2.1** (Greene-Wu). *Let  $f : M \rightarrow V$  be continuous. Let  $K$  be a compact subset of  $M$ . When  $\varepsilon > 0$  is small enough (depending only on  $M$  and  $K$ ),  $f_\varepsilon$  is a smooth function on an open neighborhood of  $K$ . The function  $f_\varepsilon$  converges to  $f$  uniformly on  $K$  as  $\varepsilon \rightarrow 0$ . If  $f$  is smooth on a neighborhood of  $K$ , then  $f_\varepsilon$  converges to  $f$  in the smooth topology on  $K$  as  $\varepsilon \rightarrow 0$ . If  $M$  is connected and  $f$  has Lipschitz constant  $D$  on a neighborhood  $W$  of  $K$ , i.e.,*

$$\sup_{x,y \in W, x \neq y} \frac{\|f(x) - f(y)\|}{\rho(x,y)} = D,$$

where  $\rho$  denotes the geodesic distance on  $M$ , then for any  $\delta > 0$ , when  $\varepsilon$  is small enough one has  $\|\partial_v(f_\varepsilon)\| < D + \delta$  for all  $x \in K$  and unit vector  $v \in TM_x$ .

One direct consequence of the above approximation theorem is

**Corollary 2.2.** *Let  $M$  is a connected closed Riemannian manifold. For any continuous map  $f : M \rightarrow V$  and any  $\delta > 0$ , when  $\varepsilon > 0$  is small enough,  $f_\varepsilon$  is a smooth function on  $M$  with  $\|f - f_\varepsilon\|_\infty < \delta$  and  $L(f_\varepsilon) < L(f) + \delta$ , where  $L(f_\varepsilon)$  and  $L(f)$  are defined by (1).*

Using Theorem 2.1 for the case  $V = \mathbb{R}$ , Greene and Wu actually showed [3, Proposition 2.1] that for any connected (possibly noncompact) Riemannian manifold  $M$  without boundary, any Lipschitz function  $f : M \rightarrow \mathbb{R}$ , any continuous function  $g : M \rightarrow \mathbb{R}_{>0}$ , and any  $\delta > 0$ , there is some smooth function  $h : M \rightarrow \mathbb{R}$  such that  $|f - h| < g$  on  $M$  and  $L(h) < L(f) + \delta$ . It would be interesting to see whether this result extends to every Banach space  $V$ .

### 3. ESTIMATE OF LIPSCHITZ CONSTANTS FOR PROJECTIONS

The key in our approach is the following estimate of the Lipschitz seminorm of the projection obtained from an ‘‘almost-projection’’ in the usual way.

**Proposition 3.1.** *Let  $A$  be a unital  $C^*$ -algebra, and let  $a \in A$  be self-adjoint whose spectrum has empty intersection with the interval  $(\delta, 1 - \delta)$  in  $\mathbb{R}$  for some  $0 < \delta < 1/2$ . Let  $L$  be a (possibly  $+\infty$ -valued) seminorm on  $A$ . Suppose that  $L$  is lower semi-continuous, vanishes on*

$1_A$ , and  $L(b^{-1}) \leq \|b^{-1}\|^2 L(b)$  for every invertible  $b \in A$ . Let  $f$  denote the characteristic function of the interval  $[1 - \delta, +\infty)$  on  $\mathbb{R}$ . Then

$$L(f(a)) \leq L(a)/(1 - 2\delta),$$

where  $f(a)$  denotes the continuous-functional calculus of  $a$  under  $f$ .

*Proof.* Take  $R > 0$  such that the spectrum of  $a$  is contained in the union of the intervals  $(-\infty, \delta]$  and  $[1 - \delta, 1 + R]$  in  $\mathbb{R}$ .

Denote by  $g$  the characteristic function of  $\{z \in \mathbb{C} : \operatorname{Re} z \geq (\delta + 1/2)/2\}$  on  $\mathbb{C}$ . Note that  $g$  is holomorphic on  $\{z \in \mathbb{C} : \operatorname{Re} z \neq (\delta + 1/2)/2\}$ . For each  $s > R$ , denote by  $\gamma_s$  the rectangle with vertices  $1/2 + si$ ,  $1/2 - si$ ,  $(1 + s) - si$ , and  $(1 + s) + si$ , parameterized as a piecewise smooth simple closed curve in the anti-clockwise direction. Then

$$\begin{aligned} L(f(a)) &= L(g(a)) = L\left(\frac{1}{2\pi i} \int_{\gamma_s} g(z)(z - a)^{-1} dz\right) \\ &\leq \frac{1}{2\pi} \int_{\gamma_s} L(g(z)(z - a)^{-1}) d|z| \\ &= \frac{1}{2\pi} \int_{\gamma_s} L((z - a)^{-1}) d|z| \\ &\leq \frac{1}{2\pi} \int_{\gamma_s} \|(z - a)^{-1}\|^2 L(z - a) d|z| \\ &= \frac{L(a)}{2\pi} \int_{\gamma_s} \|(z - a)^{-1}\|^2 d|z|, \end{aligned}$$

where the first inequality follows from the lower semi-continuity of  $L$ . For  $z$  in the line segment from  $1/2 - si$  ( $(1 + s) + si$  resp.) to  $(1 + s) - si$  ( $1/2 + si$  resp.), one has  $\|(z - a)^{-1}\|^2 \leq s^{-2}$  since  $a$  is self-adjoint. For  $z = 1/2 + ti$  with  $t \in \mathbb{R}$ , one has  $\|(z - a)^{-1}\|^2 \leq ((1/2 - \delta)^2 + t^2)^{-1}$ . For  $z = (1 + s) + ti$  with  $t \in \mathbb{R}$ , one has  $\|(z - a)^{-1}\|^2 \leq ((s - R)^2 + t^2)^{-1}$ .

Therefore

$$\begin{aligned}
L(f(a)) &\leq \frac{L(a)}{2\pi} \int_{\gamma_s} \|(z - a)^{-1}\|^2 |dz| \\
&\leq \frac{L(a)}{2\pi} (2s^{-2}(1/2 + s) + \int_{-\infty}^{+\infty} ((1/2 - \delta)^2 + t^2)^{-1} dt \\
&\quad + \int_{-\infty}^{+\infty} ((s - R)^2 + t^2)^{-1} dt) \\
&= \frac{L(a)}{2\pi} (s^{-2}(1 + 2s) + \frac{1}{1/2 - \delta} \int_{-\infty}^{+\infty} (1 + t^2)^{-1} dt \\
&\quad + \frac{1}{s - R} \int_{-\infty}^{+\infty} (1 + t^2)^{-1} dt) \\
&= \frac{L(a)}{2\pi} (s^{-2}(1 + 2s) + \frac{\pi}{1/2 - \delta} + \frac{\pi}{s - R}).
\end{aligned}$$

Letting  $s \rightarrow \infty$ , we obtain  $L(f(a)) \leq L(a)/(1 - 2\delta)$  as desired.  $\square$

Using Proposition 3.1, Rieffel has improved various estimates in a previous version of Sections 3, 4, 6 and 10 of [5].

In general the estimate in Proposition 3.1 is the best possible, as the following example shows.

**Example 3.2.** Let  $X = \{x_1, x_2\}$  be a 2-point metric space with the metric  $\rho(x_1, x_2) = 1$ . Define  $L$  on  $C(X)$  via (1). By [5, Proposition 2.2] the seminorm  $L$  satisfies the conditions in Proposition 3.1. For  $0 < \delta < 1/2$  define  $a \in C(X)$  by  $a(x_1) = \delta$  and  $a(x_2) = 1 - \delta$ . Then  $(f(a))(x_1) = 0$  and  $(f(a))(x_2) = 1$ , where  $f$  is as in Proposition 3.1. Thus  $L(a) = 1 - 2\delta$  and  $L(f(a)) = 1 = L(a)/(1 - 2\delta)$ .

We are ready to prove Theorem 1.1.

*Proof of Theorem 1.1.* Let  $0 < \delta < 1/2$ . By Corollary 2.2 applied to  $V$  equal to the self-adjoint part of  $A$  and  $f$  equal to  $p$ , we can find some self-adjoint  $p_1 \in C^\infty(M, A)$  with  $\|p - p_1\|_\infty < \delta$  and  $L(p_1) < L(p) + \varepsilon/2$ . Since  $p$  is a projection, by [5, Lemma 3.2] the spectrum of  $p_1$  is contained in the union of the intervals  $[-\delta, \delta]$  and  $[1 - \delta, 1 + \delta]$  in  $\mathbb{R}$ . Let  $f$  denote the characteristic function of the interval  $[1 - \delta, +\infty)$  on  $\mathbb{R}$ . Set  $q = f(p_1)$ .

Say,  $A$  is a real  $C^*$ -subring contained in a  $C^*$ -algebra  $B$ . Since  $C^\infty(M, B)$  is closed under the holomorphic functional calculus, we have  $q \in C^\infty(M, B)$ . Using polynomial approximations it is easy to see that the self-adjoint part of  $C(M, A)$  is closed under continuous functional calculus for real-valued continuous functions on  $\mathbb{R}$  vanishing at 0 (see for

example the proof of [5, Proposition 2.4]). Therefore  $q \in C^\infty(M, B) \cap C(M, A) = C^\infty(M, A)$ .

One has

$$\|q - p\|_\infty \leq \|q - p_1\|_\infty + \|p_1 - p\|_\infty < \delta + \delta = 2\delta.$$

It is readily checked that the Lipschitz seminorm  $L$  on the  $C^*$ -algebra  $C(M, B)$  defined by (1) satisfies the conditions in Proposition 3.1 (see for example the proof of [5, Proposition 2.2]). By Proposition 3.1 one has

$$L(q) \leq L(p_1)/(1 - 2\delta) < (L(p) + \varepsilon/2)/(1 - 2\delta).$$

Thus, when  $\delta > 0$  is small enough, we have  $\|q - p\|_\infty < \varepsilon$  and  $L(q) < L(p) + \varepsilon$  as desired.  $\square$

#### REFERENCES

- [1] R. E. Greene and H. Wu. On the subharmonicity and plurisubharmonicity of geodesically convex functions. *Indiana Univ. Math. J.* **22** (1972/73), 641–653.
- [2] R. E. Greene and H. Wu. Integrals of subharmonic functions on manifolds of nonnegative curvature. *Invent. Math.* **27** (1974), 265–298.
- [3] R. E. Greene and H. Wu.  $C^\infty$  approximations of convex, subharmonic, and plurisubharmonic functions. *Ann. Sci. École Norm. Sup. (4)* **12** (1979), no. 1, 47–84.
- [4] M. A. Rieffel. Gromov-Hausdorff distance for quantum metric spaces. *Mem. Amer. Math. Soc.* **168** (2004), no. 796, 1–65. arXiv:math.QA/0011063.
- [5] M. A. Rieffel. Vector bundles and Gromov-Hausdorff distance. *J. K-Theory* to appear. arXiv:math/0608266.

DEPARTMENT OF MATHEMATICS, SUNY AT BUFFALO, BUFFALO, NY 14260-2900, U.S.A.

*E-mail address:* hfli@math.buffalo.edu