THE CONLEY INDEX OVER A BASE

Marian Mrozek*
Instytut Informatyki
Uniwersytet Jagielloński
30-072 Kraków
Poland
mrozek@ii.uj.edu.pl

James F. Reineck
Department of Mathematics
SUNY at Buffalo
Buffalo, NY 14214-3093
USA
reineck@newton.math.buffalo.edu

Roman Srzednicki†
Instytut Matematyki
Uniwersytet Jagielloński
30-059 Kraków
Poland
srzednic@im.uj.edu.pl

August 19, 1996

Abstract

We construct a generalization of the Conley index for flows. The new index preserves information which in the classical case is lost in the process of collapsing the exit set to a point. The new index has most of the properties of the classical index. As examples, we study a flow with a knotted orbit in R^3, and the problem of continuing two periodic orbits which are not homotopic as loops.

Classifications: Primary 34C35; Secondary 58F25, 34A34

Keywords: Conley index, continuation, fiberwise pointed space

1 Introduction

The Conley index is a useful tool in the study of flows. To obtain the index of an isolated invariant set S, one takes an index pair (P_1, P_2) for S and collapses P_2 to a point. (Recall that P_1 is an isolating neighborhood for S, and P_2 is an exit set for P_1.) The homotopy type of the resulting quotient space is independent of the choice of index pair. Indeed, given two index pairs, after collapsing the exit sets there is a flow-defined homotopy equivalence between the resulting quotient spaces. The homotopy type of the quotient space, or more precisely, the collection of all such quotients and homotopy equivalences is called the Conley index of S, and is denoted $h(S)$.

One weakness of the index is that information may be lost when P_2 is collapsed to a point. An example from Conley's monograph [Co] illustrates this.

In Figure 1, P_1 is a cylinder from which a tubular neighborhood of a knotted orbit has been removed, and P_2 is the punctured disc at the bottom of P_1. P_2 is not a strong deformation retract of P_1, so by Ważewski's theorem, there is a point whose forward orbit is contained in P_1, and the omega-limit set of this orbit is an invariant set contained in P_1. However, when P_2 is collapsed to a point, the resulting quotient space is homotopic to a point, and thus the Conley index does not detect the invariant set. Indeed, a

*Supported by KBN, Grant 0449/P3/94/06.
†Supported by KBN, Grant 2 P03A 040 10.



Figure 1: A knotted orbit in R^3.

question posed in [Co] asks how much information is lost under this collapse, and how it can be retained.

The Conley index is a continuation invariant. If one has an isolating neighborhood N, and deforms the flow in such a way that N remains an isolating neighborhood throughout the deformation, then the sets isolated by N are related by continuation, and have the same index. An obvious question is the converse: if two sets have the same index, are they related by continuation? It is easy to find examples to show that this need not be the case. For example, if the phase space is disconnected, it is easy to construct flows with critical points of the same index in different components, and these sets are not related by continuation. The problem here is that the index does not take into account how the set lies in the phase space. In the example with a disconnected phase space, it is obvious that there is no continuation, but a more subtle question is the following: can two periodic orbits which are not homotopic as loops be related by continuation? A repelling periodic orbit in the punctured plane has the same index as the disjoint union of two critical points, so it is not so clear that there is no continuation between periodic orbits in different homotopy classes.

In this paper, we generalize the index by gluing the exit set P_2 of a regular index pair to a topological space Z, which we call the base space. The gluing map $\omega : X \to Z$ is defined on the phase space X, and the index of a set S over Z is the weak fiberwise homotopy type of the adjunction space $P_1 \cup_{\omega|_{P_2}} Z$. The precise definition of weak fiberwise homotopy type is given below, but

it is a finer invariant than the usual homotopy type of the adjunction, as Example 8.3 illustrates. If Z is a point, then this gives the classical Conley index. Another possibility for an arbitrary flow is $Z = X$, $\omega = \mathrm{id}$. These two examples represent the maximal and minimal information loss due to collapsing. Other spaces Z and maps ω lie somewhere in between in terms of information lost upon collapse.

The index over a base has properties similar to the classical Conley index. The homotopy type (more precisely the weak fiberwise homotopy type) of the quotient $P_1 \cup_{\omega|_{P_2}} Z$ is independent of the choice of regular index pair, and is invariant under continuation. The construction of the index over a base is fully functorial. In particular, this means that the new index extends to a connected simple system, and one may also obtain a Conley functor as in [KM]. By making a judicious choice of Z and ω, one can often obtain more information than in the classical case. For example, in the flow from Figure 1, if one chooses Z to be P_1, and ω to be the identity, then the space $P_1 \cup_{\omega|_{P_2}} Z$ is not homotopically equivalent to Z, so the maximal invariant set in P_1 is nonempty. Thus an invariant set which has a trivial classical Conley index may have a nontrivial index over an appropriate base.

In the case of periodic orbits in different homotopy classes, let X denote the punctured plane $\mathrm{R}^2 \setminus \{0\}$, Z be the unit circle, and $\omega(x) = x/|x|$. Suppose S is a repelling periodic orbit, with P_1 an annulus, and $P_2 = \partial P_1$. If S is homotopically nontrivial, then the index of S over Z is a 2-torus, but if S is homotopically trivial, it is the wedge of a 2-sphere and two 1-spheres. Thus these sets are not related by continuation in any family of flows on the punctured plane.

We were originally led to this generalization of the index by studying the classical Conley index for periodic orbits. In the case of hyperbolic periodic orbits, our generalized index has a structure similar to that of a bundle over the circle as in the noncontractible orbit in the punctured plane. Bundles over the circle are classified by a single gluing map on the fiber. In the periodic orbit case that map is related to the Poincaré map. In particular, all information carried by the Conley index of the Poincaré map may be regained from an appropriate Conley index over S^1. This result will be described precisely in a future paper.

In this paper, our setting is the case of semiflows on locally compact metric spaces, but the same ideas can be applied to infinite dimensional semiflows such as those generated by partial differential equations in the setting studied

by Rybakowski (see [Ry]). Since the additional technical details needed in the infinite dimensional case would obscure the main ideas of this paper, we only deal with the finite dimensional case. However we work with semiflows, and all the topological background is developed to work in the general case; in particular, in Section 4 we deal with pairs of closed subspaces instead of compact ones. Thus the reader familiar with the Rybakowski approach will have no problem extending the theory presented here to the Rybakowski setting.

There is a paper with a similar title by Bartsch [Ba], but the similarity does not go beyond the title. We cite his paper to avoid confusion.

We denote the sets of all reals, integers, nonnegative integers, and nonpositive integers by $\mathrm{R}, \mathrm{Z}, \mathrm{Z}^+$, and Z^-, respectively. For a given topological space X and $A \subset X$, we denote the closure of A by $\operatorname{cl} A$, its interior by $\operatorname{int} A$ and its boundary by $\operatorname{bd} A$.

Acknowledgement. We thank Professor F. W. Lawvere for informing us about the literature on fiberwise pointed spaces.

2 Main Results

In this section we state our main results. We assume the reader is familiar with the standard concepts of Conley index theory as presented in [Co, Sm]. Precise definitions are given later. Let X be a locally compact metric space and let φ be a semiflow on X. For an index pair $P = (P_1, P_2)$ in a neighborhood of some isolated invariant set, define the map $\tau_P : P_1 \to [0, \infty]$ by

$$\tau_P(x) := \begin{cases} \sup\{t \geq 0 \mid x \cdot [0,t] \subset P_1 \setminus P_2\} & \text{if } x \in P_1 \setminus P_2 \\ 0 & \text{if } x \in P_2. \end{cases}$$

Recall that an index pair in a flow is called *regular* if τ_P is continuous. In the case of semiflows the definition is somewhat more complicated, and is given in Section 5.

Let Z be another Hausdorff space and $\omega : X \to Z$ be a continuous map. For the regular index pair $P = (P_1, P_2)$ we define $U_\omega(P)$ as the adjunction $P_1 \cup_{\omega|_{P_2}} Z$, i.e.

$$U_\omega(P) := Z \times 0 \cup P_1 \times 1/\sim,$$

where $\sim$ denotes the minimal equivalence relation which identifies a point $(x,1) \in P_2 \times 1$ with $(\omega(x),0) \in Z \times 0$. Denote by $[u,i]$ the equivalence class of (u,i) in $U_\omega(P)$. There are a natural embedding $s_P : Z \to U_\omega(P)$, $z \mapsto [z,0]$ and a natural projection $r_P : U_\omega(P) \to Z$; $r_P([x,1]) = \omega(x)$ and $r_P([z,0]) = z$. Obviously, $r_P \circ s_P = \mathrm{id}_Z$.

As in the case of the classical Conley index theory, the homotopy type of $U_\omega(P)$ is independent of the choice of (regular) index pair. In fact, a stronger statement is true. In order to state it, we introduce the notion of weak homotopy type. Let (U,r,s) be a triple consiting of a topological space U and continuous maps $r : U \to Z$ and $s : Z \to U$ such that $r \circ s = \mathrm{id}_Z$. We call such a triple a *fiberwise pointed space* (over Z). If Z is a one-point space then this notion coincides with the notion of a topological space with a base-point. As we have shown above, $(U_\omega(P), r_P, s_P)$ is a fiberwise pointed space. Two fiberwise pointed spaces (U,r,s) and (U',r',s') are equivalent if there exist continuous maps $f : U \to U'$ and $f' : U' \to U$ satisfying

$$f \circ s = s', \quad f' \circ s' = s,$$

$$r' \circ f \simeq r \operatorname{rel} s(Z), \quad r \circ f' \simeq r' \operatorname{rel} s'(Z),$$

$$f' \circ f \simeq \mathrm{id}_U \operatorname{rel} s(Z), \quad f \circ f' \simeq \mathrm{id}_{U'} \operatorname{rel} s'(Z).$$

The equivalence class is called the *weak homotopy type over* Z. The type of $(Z, \mathrm{id}_Z, \mathrm{id}_Z)$ is called *trivial*. The wedge sum of weak homotopy types is defined in an obvious way. On the other hand the smash product is not defined in general; in order to define it correctly one should consider *fiberwise well-pointed spaces over* Z, i.e. fiberwise pointed spaces (U,r,s) such that s is a cofibration. Actually, if P is a regular index pair then $(U_\omega(P), r_P, s_P)$ is well-pointed.

Theorem 2.1 *If P and Q are two regular index pairs for an isolated invariant set S then $(U_\omega(P), r_P, s_P)$ and $(U_\omega(Q), r_Q, s_Q)$ have the same weak homotopy type over Z.*

The weak homotopy type over Z described in Theorem 2.1 is denoted by $h_\omega(S,\varphi)$ and is called the *Conley index of S over the base Z*. Of course, the index also depends on the map ω (actually: on the homotopy class of ω), hence the notation. If ω is a constant map, then the resulting index is simply the wedge of the classical Conley index and Z.

This generalized index has many of the properties of the classical index. Clearly, if $h_\omega(S,\varphi)$ is not trivial then S is nonempty. The index of the disjoint union of two isolated invariant sets is the wedge of the indices and the index of the cartesian product of two isolated invariant sets is the smash product of their indices. As in the classical case, the index over Z is invariant under continuation. Consider a family of semiflows $\varphi^\lambda : X \times \mathrm{R}^+ \to X$ continuously depending on $\lambda \in [0,1]$. We have the following theorem.

Theorem 2.2 *If N is an isolating neighborhood with respect to φ^λ for all $\lambda \in [0,1]$ then $h_\omega(N,\varphi^\lambda)$ does not depend on $\lambda \in \Lambda$.*

Theorems 2.1 and 2.2 will be proved in sections 6 and 7 respectively.

3 Fiberwise pointed spaces over a base

In the sequel Z denotes a fixed topological space. A triple $\mathbf{U} = (U, r, s)$ is called a *fiberwise pointed (topological) space over Z* if U is a topological space, $r : U \to Z$ and $s : Z \to U$ are continuous maps, and $r \circ s = \mathrm{id}_Z$ (compare [J3, p. 41–42]). In [J1, p. 163] and [J2, p. 98] such a space is also called a sectioned space over Z. We refer to Z as the base space, to r as the projection, and to s as the section of $\mathbf{U}$. The projection is an r-map in the sense of [Bo]. In particular it follows that $s(Z)$ is a retract (hence a closed subset) of U, $s \circ r : U \to s(Z)$ is a retraction, and s and the restriction of r are mutually inverse homeomorphisms between Z and $s(Z)$.

We want to turn the fiberwise pointed spaces into a category. Let $\mathbf{U}' = (U', r', s')$ be another fiberwise pointed space over Z. A natural choice for a morphism from $\mathbf{U}$ to $\mathbf{U}'$ is to take a *fiberwise pointed* map, i.e. a continuous map $\phi : U \to U'$ such that $r' \circ \phi = r$ and $\phi \circ s = s'$ (comp. [J1, J3] for instance). For our purposes such a definition is too strong.

We define a *homotopically fiberwise pointed map* (shortly: an *h.f.p. map*) $f : \mathbf{U} \to \mathbf{U}'$ as a continuous map $f : (U, s(Z)) \to (U', s'(Z))$ such that

$$r' \circ f \simeq r \operatorname{rel} s(Z).$$

Obviously a fiberwise pointed map is an h.f.p. map. Note that there always exists a fiberwise pointed map (and consequently an h.f.p. map) between $\mathbf{U}$ and $\mathbf{U}'$, namely $s' \circ r$ (it is called the *fiberwise constant map*).

Proposition 3.1 *If $f : \mathbf{U} \to \mathbf{U}'$ is h.f.p., then $f \circ s = s'$ and the restriction of f to a map $s(Z) \to s'(Z)$ is equal to the homeomorphism $s' \circ r|_{s(Z)}$. In particular $f(s(Z)) = s'(Z)$.* $\square$

One can easily verify that the identity map is an h.f.p. map and the composition of h.f.p. maps is an h.f.p. map. Thus fiberwise pointed spaces over Z together with h.f.p. maps constitute a category. We will call it the weak category of fiberwise pointed topological spaces and we will denote it by Fib_Z.

We define $\mathbf{U} \vee \mathbf{U}'$, the *fiberwise pointed wedge*, as the triple $(U \vee U', r_\vee, s_\vee)$ such that $U \vee U'$ is the disjoint union of U and U' with $s(z)$ and $s'(z)$ identified, i.e. $U \vee U' := U \times 0 \cup U' \times 1/\sim_\vee$, where $\sim_\vee$ denotes the equivalence relation generated by $(s(z), 0) \sim_\vee (s'(z), 1)$ for any $z \in Z$. The maps $r_\vee, s_\vee$ are given by

$$r_\vee([(x,i)]) := \begin{cases} r(x) & \text{if } x \in U \\ r'(x) & \text{if } x \in U'. \end{cases}$$

and $s_\vee(z) := [(s(z), 0)] = [(s'(z), 0)]$. It is easy to check that $r_\vee$ is well defined, that $\mathbf{U} \vee \mathbf{U}'$ is a fiberwise pointed space, and that this agrees with the usual definition of wedge when Z is a single point.

Let $\mathbf{U} = (U, r, s)$ and $\mathbf{U}' = (U', r', s')$ be two fiberwise pointed spaces over Z and Z', respectively. In the cartesian product $U \times U'$ define an equivalence relation $\sim_\wedge$ by $(u, u') \sim_\wedge (v, v')$ if $(u, u') = (v, v')$ or $u = v \in s(Z)$, $r'(u') = r'(v')$ or $r(u) = r(v)$, $u' = v' \in s'(Z')$. Put

$$U \wedge U' := U \times U'/\sim_\wedge$$

and denote by $u \wedge u'$ the equivalence class of (u, u'). Define maps $r_\wedge : U \wedge U' \to Z \times Z'$ and $s_\wedge : Z \times Z' \to U \wedge U'$ by

$$r_\wedge(u \wedge u') := (r(u), r'(u')), \quad s_\wedge(z, z') := s(z) \wedge s'(z').$$

The triple

$$\mathbf{U} \wedge \mathbf{U}' := (U \wedge U', r_\wedge, s_\wedge)$$

is a fiberwise pointed space over $Z \times Z'$ and is called the *fiberwise pointed smash product* of $\mathbf{U}$ and $\mathbf{U}'$. Again, if both Z and Z' are one-pointed spaces than this notion agrees with the usual smash product.

By a *homotopy* in the category Fib_Z we mean a collection of h.f.p. maps $F_t : \mathbf{U} \to \mathbf{U}'$, $t \in [0,1]$ such that the map $F : U \times [0,1] \ni (x,t) \to F_t(x) \in U'$ is continuous. We say that two h.f.p. maps $f, g : \mathbf{U} \to \mathbf{U}'$ are homotopic in Fib_Z if there exists a homotopy $F_t : \mathbf{U} \to \mathbf{U}'$, $t \in [0,1]$ in Fib_Z such that $F_0 = f, F_1 = g$.

As an easy consequence of Proposition 3.1 we obtain the following

Proposition 3.2 *Assume $f : \mathbf{U} \to \mathbf{U}'$ is an h.f.p. map and $g : U \to U'$ is continuous. If*

$$f \simeq g \operatorname{rel} s(Z)$$

then g is also an h.f.p. map from $\mathbf{U}$ to $\mathbf{U}'$ and f and g are homotopic in Fib_Z.

Proof: We have $f \simeq g \operatorname{rel} s(Z)$, thus also $r'f \simeq r'g \operatorname{rel} s(Z)$. But f is a morphism in Fib_Z, hence $r \simeq r'f \simeq r'g \operatorname{rel} s(Z)$. This shows that also g is a morphism in Fib_Z. Let F_t be the homotopy joining f to g. Then $F_t \simeq f \operatorname{rel} s(Z)$ for each $t \in [0,1]$. It follows from what we already proved that F_t is an h.f.p. map and consequently f and g are homotopic in Fib_Z. □

Obviously, the homotopy in the category Fib_Z induces an equivalence relation between the h.f.p. maps. Taking equivalence classes of h.f.p. maps as a new collection of morphisms we obtain another category, which we will call the homotopy weak category of fiberwise pointed spaces. We will denote it by HtpFib_Z. Obviously we have also the homotopy functor $\mathrm{Htp} : \mathrm{Fib}_Z \to \mathrm{HtpFib}_Z$ sending each h.f.p. map to its equivalence class. An h.f.p. map ϕ will be called a *weak homotopy equivalence* if $\mathrm{Htp}(\phi)$ is an isomorphism in HtpFib_Z.

Proposition 3.3 *Assume $f : \mathbf{U} \to \mathbf{U}'$ is an h.f.p. map. If there exists a continuous map $f' : U' \to U$ such that*

$$f' \circ f \simeq \mathrm{id}_U \operatorname{rel} s(Z), \quad f \circ f' \simeq \mathrm{id}_{U'} \operatorname{rel} s'(Z).$$

then f' is also an h.f.p. map and f and f' are mutually inverse weak homotopy equivalences.

Proof: First observe that $f'f \simeq \mathrm{id}_U \operatorname{rel} s(Z)$ implies that $f'f(s(Z)) = s(Z)$. From the homotopy equivalence $ff' \simeq \mathrm{id}_U \operatorname{rel} s(Z)$ we get $r'ff' \simeq r' \operatorname{rel} s(Z')$

and $rf' \simeq r' \mathrm{rel} s'(Z)$, because f is a morphism in Fib_Z. It follows that f' is also a morphism in Fib_Z and consequently f and f' are mutually inverse weak homotopy equivalences. $\square$

We say that two fiberwise pointed spaces $\mathbf{U}$ and $\mathbf{U}'$ over Z have the same *weak fiberwise pointed homotopy type over* Z (shortly: the weak homotopy type) if there exists a weak homotopy equivalence between them. (It should be pointed out that we use the term "weak homotopy type" in a different meaning then it is commonly used in topology, compare for example [Wh].) The equivalence class of spaces with the same weak homotopy type as $\mathbf{U}$ is denoted by $[\mathbf{U}]_Z$. A distinguished role is played by

$$\overline{0}_Z := [(Z, \mathrm{id}_Z, \mathrm{id}_Z)]_Z,$$

the *trivial* weak homotopy type over Z.

We compare various homotopy types on a simple example.

Example 3.4 For $x \in \mathrm{R} \setminus 0$ put $V_x = (\mathrm{R} \setminus 0) \times 0 \cup \{x\} \times 1$, i.e. V_x is the disjoint union of $\mathrm{R} \setminus 0$ and $\{x\}$. Define $r_x : V_x \to \mathrm{R} \setminus 0$ and $s_x : \mathrm{R} \setminus 0 \to V_x$ by $r_x(y,i) = y$ and $s_x(z) = (z,0)$. Then $\mathbf{V}_x = (V_x, r_x, s_x)$ is a fiberwise pointed space over $\mathrm{R} \setminus 0$. Each of the spaces V_x has the same (ordinary) homotopy type of a three-point space. It is easy to see, that for $x \neq x'$ there is no fiberwise pointed map $\mathbf{V}_x \to \mathbf{V}_{x'}$ except of the constant one, hence the strong homotopy types (i.e. the homotopy types in the sense of [J1, J3]) of $\mathbf{V}_x$ are all different. On the other hand there are no nonconstant h.f.p. maps between $\mathbf{V}_x$ and $\mathbf{V}_{x'}$ only in the case where the signs of x and x' are opposite, hence there are exactly two different weak homotopy types, and $[\mathbf{V}_x]_{\mathrm{R}\setminus 0} = [\mathbf{V}_{x'}]_{\mathrm{R}\setminus 0}$ iff $xx' > 0$.

In the above example we compared the weak homotopy types of spaces with several connected components. This can be generalized to the following simple observation.

Proposition 3.5 *Let $\{T_i\}_{i=0,\dots,r}$ and $\{T'_i\}_{i=0,\dots,r}$ be the sets of path components of U and U', respectively. Let*

$$s(Z) \subset T_0, \ s'(Z) \subset T'_0.$$

If $f : \mathbf{U} \to \mathbf{U}'$ is a weak homotopy equivalence then there is a permutation σ, $\sigma(0) = 0$, such that for $i = 0, \dots, r$: $f(T_i) \subset T'_{\sigma(i)}$ and the restriction $f_i : T_i \to T_{\sigma(i)}$ of f is a homotopy equivalence such that $r' \circ f_i \simeq r|_{T_i}$. $\square$

In order to distinguish weak homotopy types of $\mathbf{U}$ and $\mathbf{U}'$ one should compare the homotopy types $[U]$ and $[U']$ first. If they are equal, further necessary information can be provided by Proposition 3.5 as the next example shows:

Example 3.6 Let $W_1 = W_2 = S^1 \times 0 \cup S^1 \times 1$, i.e. W_1 and W_2 are equal to the disjoint union of two copies of the circle $S^1 = \{x \in \mathrm{C} : |x| = 1\}$. Define $r_i : W_i \to S^1$ and $s_i : S^1 \to W_i$, $i = 1, 2$ by

$$r_1(x, i) = x,\ s_1(y) = (y, 0),$$

$$r_2(x, 0) = x,\ r_2(x, 1) = x^2,\ s_2(y) = (y, 0).$$

Then $\mathbf{W}_i = (W_i, r_i, s_i)$, $i = 1, 2$ are fiberwise pointed spaces over S^1. It follows by Proposition 3.5 that if $f : \mathbf{W}_1 \to \mathbf{W}_2$ is a weak homotopy equivalence then $f(S^1 \times 1) \subset (S^1 \times 1)$ and $r_2 \circ f|_{S^1 \times 1} \simeq r_1|_{S^1 \times 1}$, which is impossible. Thus $[\mathbf{W}_1]_{S^1} \neq [\mathbf{W}_2]_{S^1}$.

Remark 3.7 If we consider the question of comparison of $[\mathbf{U}]_Z$ and $[\mathbf{U}']_Z$ on the homology level, one would expect that the information provided by the triples $(H(U), H(r), H(s))$ and $(H(U'), H(r'), H(s'))$ is essentially better than the one provided by $H(U)$ and $H(U')$ alone. However, this is not the case, at least in reasonable situations: if $H(U) \cong H(U')$ and $H(Z)$ is a finitely generated abelian group then the diagram



is commutative for some isomorphism ϕ.

Indeed, let $j : U \hookrightarrow (U, s(Z))$ and $j' : U' \hookrightarrow (U', s'(Z))$ be the inclusions. Since $s(Z)$ is a retract of U, the exact sequence

$$0 \longrightarrow H(Z) \xrightarrow{H(s)} H(U) \xrightarrow{H(j)} H(U, s(Z)) \longrightarrow 0$$

splits and $(H(r), H(j)) : H(U) \to H(Z) \oplus H(U, s(Z))$ is an isomorphism. Up to this isomorphism $H(r)$ and $H(s)$ are represented as the maps

$$H(Z) \oplus H(U, s(Z)) \ni (u, v) \longrightarrow u \in H(Z),$$

$$H(Z) \ni u \longrightarrow (u, 0) \in H(Z) \oplus H(U, s(Z)),$$

respectively. The same holds true for the triple $(H(U'), H(r'), H(s'))$. It follows by the assumption on $H(Z)$ and by [Coh] or [W] that there exists an isomorphism $\psi : H(U, s(Z)) \to H(U', s'(Z))$, and we can put

$$\phi = (H(r'), H(j'))^{-1} \circ (\mathrm{id}_{H(Z)} \oplus \psi) \circ (H(r), H(j)).$$

It is easily seen that if $[\mathbf{U}]_Z = [\mathbf{U}']_Z$ and $[\mathbf{V}]_Z = [\mathbf{V}']_Z$ then $[\mathbf{U} \vee \mathbf{V}]_Z = [\mathbf{U}' \vee \mathbf{V}']_Z$, hence we can define

$$[\mathbf{U}]_Z \vee [\mathbf{V}]_Z := [\mathbf{U} \vee \mathbf{V}]_Z,$$

the wedge of weak homotopy types. Obviously, for every space $\mathbf{U}$,

$$[\mathbf{U}]_Z \vee \overline{0}_Z = \overline{0}_Z \vee [\mathbf{U}]_Z = [\mathbf{U}]_Z.$$

On the other hand, we cannot provide a general definition of smash product of week homotopy types, as the following example shows:

Example 3.8 Let

$$U = I \times 0 \cup \{(0, 1)\} \subset \mathrm{R}^2, \quad U' = I \times 0 \cup \{(1, 1)\} \subset \mathrm{R}^2.$$

Assume that both $r : U \to I$ and $r' : U' \to I$ are given by $(x, y) \mapsto x$, and both $s : I \to U$ and $s' : I \to U'$ are given by $x \mapsto (x, 0)$. Let

$$V = \{\frac{1}{n} : n \in \mathrm{Z}^+\} \cup \{0\} \subset \mathrm{R},$$

$\rho : V \to \{0\}$ and $\sigma : \{0\} \to V$, $\sigma(0) = 0$. Put $\mathbf{U} = (U, r, s)$, $\mathbf{U}' = (U', r', s')$, and $\mathbf{V} = (V, \rho, \sigma)$. One can show that $[\mathbf{U}]_I = [\mathbf{U}']_I$, but

$$[\mathbf{U} \wedge \mathbf{V}]_{I \times 0} \neq [\mathbf{U}' \wedge \mathbf{V}]_{I \times 0}.$$

In order to verify this assertion we introduce new, simpler spaces

$$W = I \times 0 \cup \{(0, \frac{1}{n}) : n \in \mathrm{Z}^+\}, \quad W' = I \times 0 \cup \{(1, \frac{1}{n}) : n \in \mathrm{Z}^+\},$$

mappings $R : W \to I$, $(x, y) \mapsto x$, $S : I \to W$, $x \mapsto (x, 0)$, and analogous mappings R' and S' for W'. Put $\mathbf{W} = (W, R, S)$ and $\mathbf{W}' = (W', R', S')$. The map $U \wedge V \to W$ given by $[((0,1), z)] \mapsto (0, z)$ and $[((x, 0)), z)] \mapsto (x, 0)$ establishes the isomorphism (up to the identification $I = I \times 0$) between $\mathbf{U} \wedge \mathbf{V}$ and $\mathbf{W}$. Similarly, there is an isomorphism between $\mathbf{U}' \wedge \mathbf{V}$ and $\mathbf{W}'$, hence it suffices to prove that $[\mathbf{W}]_I \neq [\mathbf{W}']_I$.

Let $f : \mathbf{W} \to \mathbf{W}'$ be an h.f.p. map. Then there is an $n_0 \neq 0$ such that $f(0, 1/n_0) \in I \times 0 \subset W'$ because $f(0,0) = (0,0)$. If $f : \mathbf{W}' \to \mathbf{W}$ is any h.f.p. map then $(f' \circ f)(0, 1/n_0) = f(0, 1/n_0)$ is contained in $I \times 0$, hence in a different path component then $\{(0, 1/n_0)\}$ and thus $f' \circ f$ cannot be homotopic to the identity, hence the types must be different.

By the previous example it is clear that in order to define the smash product one should introduce a more restrictive class of spaces than fiberwise pointed. Recall that the pair of spaces (X, A), with A closed, is called *cofiberd* if $X \times 0 \cup A \times I$ is a retract of $X \times I$ or, equivalently, if (X, A) admits a *Strøm structure*, i.e. a pair (G, a) of continuous maps $a : X \to I$ and $G : X \times I \ni (x, t) \to G_t(x) \in X$ such that $a^{-1}(0) = A$ and

$$G_0 = \mathrm{id}_X, \quad G_t(x) = x \text{ if } (x, t) \in A \times I, \quad G_1(x) \in A \text{ if } a(x) < 1$$

(compare [J2, p. 168, 169, 172] and [Wh, p. 22, 26]; in the latter reference the term "NDR-pair" is used).

We call (U, r, s) a *fiberwise well-pointed space over* Z provided it is fiberwise pointed and $(U, s(Z))$ is a cofiberd pair (or equivalently: s is a cofibration). In [J2, p. 180] such a space is called well-sectioned over Z.

Remark 3.9 In proofs of some results presented below the following fact will be useful: If $\sim$ is an equivalence relation on a topological space X and $H : X \times I \to Y$ is continuous such that $H(x, t) = H(x', t)$ if $x \sim x'$, then the induced map $H' : (X/\sim) \times I \to Y$ given by $H'([x]_\sim, t) = H(x, t)$, is continuous (compare [Sw, 0.8]).

Proposition 3.10 *Assume that* $\mathbf{U}$, $\mathbf{U}'$, $\mathbf{V}$, *and* $\mathbf{V}'$ *are fiberwise well-pointed spaces,* $\mathbf{U}$ *and* $\mathbf{U}'$ *over* Z, $\mathbf{V}$ *and* $\mathbf{V}'$ *over* Ω. *If* $[\mathbf{U}]_Z = [\mathbf{U}']_Z$ *and* $[\mathbf{U}]_\Omega = [\mathbf{U}']_\Omega$ *then*

$$[\mathbf{U} \wedge \mathbf{V}]_{Z\times\Omega} = [\mathbf{U}' \wedge \mathbf{V}']_{Z\times\Omega}.$$

Proof: In order to simplify notation, we assume (without loss of generality) that $\mathbf{V} = \mathbf{V}'$. Let $\mathbf{U} = (U, r, s)$, $\mathbf{U}' = (U', r', s')$, and $\mathbf{V} = (V, \rho, \sigma)$. Let $f : \mathbf{U} \to \mathbf{U}'$ and $f' : \mathbf{U}' \to \mathbf{U}$ be mutually inverse homotopy equivalences in Fib_Z. At first observe that one cannot usually factorize the map $(u, v) \mapsto f(u)\wedge v$ because $r(u) = r(u_1)$ does not imply $r'f(u) = r'f(u_1)$ in general, hence the definition of a homotopy equivalence from $\mathbf{U} \wedge \mathbf{V}$ to $\mathbf{U}' \wedge \mathbf{V}$ must be essentially more complicated.

Denote by $h : U \times I \to Z$ a homotopy joining r with $r' \circ f$, i.e. $h_0 = r$, $h_1 = r'f$, and $h_t s = \mathrm{id}_Z$ for $t \in I$. Let h' be an analogous homotopy joining r' with $r \circ f'$. Let (G, a), (G', a'), and (Γ, α) be Strøm structures for $(U, s(Z))$, $(U', s'(Z))$, and $(V, \sigma(Z))$, respectively. For $u \in U$, $u' \in U'$, and $v \in V$ put

$$m(u, v) = \min\{a(u), \alpha(v)\}, \quad m'(u', v) = \min\{a'(u'), \alpha(v)\}.$$

For $t \in (0, 1]$ define maps $F_t : \mathbf{U} \wedge \mathbf{V} \to \mathbf{U}' \wedge \mathbf{V}$ and $F'_t : \mathbf{U}' \wedge \mathbf{V} \to \mathbf{U} \wedge \mathbf{V}$ by

$$F_t(u\wedge v) := \begin{cases} s'h_{\frac{4}{t}m(u,v)}G_{\frac{4}{t}m(u,v)}(u)\wedge\Gamma_{\frac{4}{t}m(u,v)}(v) & \text{if } u \in \mathrm{cl}\, a^{-1}([0, \frac{1}{4}t)) \\ & \text{or } v \in \mathrm{cl}\,\alpha^{-1}([0, \frac{1}{4}t)), \\ fG_{2-\frac{4}{t}m(u,v)}(u)\wedge\Gamma_{2-\frac{4}{t}m(u,v)}(v) & \text{if } \frac{1}{4}t \le m(u, v) \le \frac{1}{2}t, \\ f(u)\wedge v & \text{if } \frac{1}{2}t \le m(u, v), \end{cases}$$

$$F'_t(u'\wedge v) := \begin{cases} sh'_{\frac{4}{t}m'(u',v)}G'_{\frac{4}{t}m'(u',v)}(u')\wedge\Gamma_{\frac{4}{t}m'(u',v)}(v) & \text{if } u' \in \mathrm{cl}(a')^{-1}([0, \frac{1}{4}t)) \\ & \text{or } v \in \mathrm{cl}\,\alpha^{-1}([0, \frac{1}{4}t)), \\ f'G'_{2-\frac{4}{t}m'(u',v)}(u')\wedge\Gamma_{2-\frac{4}{t}m'(u',v)}(v) & \text{if } \frac{1}{4}t \le m'(u', v) \le \frac{1}{2}t, \\ f'(u')\wedge v & \text{if } \frac{1}{2}t \le m'(u', v). \end{cases}$$

By the definitions of smash product and Strøm structure one can verify that the above formulas give correctly defined continuous maps which are h.f.p.

We will prove that F_1 and F_1' are mutually inverse homotopy equivalences. For $t \in (0,1]$ put

$$Y_t = \{u \wedge v \in U \wedge V : m(u,v) \geq \frac{1}{2}t\}.$$

For $u \wedge v \in Y_t$ and $\tau \in I$ put

$$n(u,v,\tau) := m'(f(u),v) + 2\tau(m(u,v) - \frac{1}{2}t)$$

and define maps $K_{t,\tau} : Y_t \to U' \wedge V$ by

$$K_{t,\tau}(u \wedge v) :=$$
$$\begin{cases} sh'_{\frac{4}{t}n(u,v,\tau)} G'_{\frac{4}{t}n(u,v,\tau)} f(u) \wedge \Gamma_{\frac{4}{t}n(u,v,\tau)}(v) \\ \quad \text{if } f(u) \in \mathrm{cl}(a')^{-1}((-\infty, \frac{1}{4}t - 2\tau(m(u,v) - \frac{1}{2}t)), \\ f' G'_{2-\frac{4}{t}n(u,v,\tau)} f(u) \wedge \Gamma_{2-\frac{4}{t}n(u,v,\tau)}(v) \\ \quad \text{if } m'(f(u),v) \in [\frac{1}{4}t - 2\tau(m(u,v) - \frac{1}{2}t), \frac{1}{2}t - 2\tau(m(u,v) - \frac{1}{2}t)], \\ f'f(u) \wedge v \text{ if } m'(f(u),v) \in [\frac{1}{2}t - 2\tau(m(u,v) - \frac{1}{2}t), \infty). \end{cases}$$

It follows in particular that $K_{t,0} = F_t' \circ F_t|_{Y_t}$, if $m(u,v) = \frac{1}{2}t$ then $K_{t,\tau}(u \wedge v) = F_t' F_t(u \wedge v)$ and if $m(u,v) \geq \frac{3}{4}t$ then $K_{t,1}(u \wedge v) = f'f(u) \wedge v$. Let $D : U \times I \to U$, $(u,t) \mapsto D_t(u)$ be the homotopy relative to $s(Z)$ joining $f'f$ with the identity, i.e.

$$D_0 = f' \circ f, \quad D_1 = \mathrm{id}_U .$$

For $t \in (0,1]$ and $\tau \in I$ define maps $H_{t,\tau} : U \wedge V \to U' \wedge V$ by

$$H_{t,\tau}(u \wedge v) := \begin{cases} F_t' F_t(u \wedge v) & \text{if } m(u,v) \leq \frac{1}{2}t, \\ K_{t,1}(u \wedge v) & \text{if } \frac{1}{2} \leq m(u,v) \leq \frac{3}{4}t, \\ D_{\frac{4\tau}{t}m(u,v) - 3\tau}(u) \wedge v & \text{if } \frac{3}{4} \leq m(u,v) \leq t, \\ D_\tau(u) \wedge v & \text{if } t \leq m(u,v). \end{cases}$$

It follows that $F_1' \circ F_1 \simeq H_{1,0} \operatorname{rel} s_\wedge(Z \times \Omega)$ via the homotopy $\{F_1' \circ F_1|_{Y_1} \cup K_{1,t}\}_{t \in I}$ and $H_{1,0} \simeq H_{1,1} \operatorname{rel} s_\wedge(Z \times \Omega)$ via the homotopy $\{H_{1,t}\}_{t \in I}$. Define the final homotopy $J : (U \wedge V) \times I \ni (u \wedge v, t) \to J_t(u \wedge v) \in U' \wedge V$ by

$$J_t := \begin{cases} \mathrm{id}_{U \wedge V} & \text{if } t = 0, \\ H_{t,1} & \text{if } t > 0. \end{cases}$$

Observe, that J is continuous in $\{(u\wedge v,0) : u \in s(Z) \text{ or } \sigma(\Omega)\}$ (hence continuous everywhere) and it is a homotopy relative to $s_\wedge(Z\times\Omega)$ joining $H_{1,1}$ with $\mathrm{id}_{U\wedge V}$. By composing the above homotopies we conclude that $F_1'\circ F_1$ is homotopic to the identity in $\mathrm{Fib}_{Z\times\Omega}$. By the symmeric argument one can show that $F_1\circ F_1' \simeq \mathrm{id}_{U'\wedge V}$ in $\mathrm{Fib}_{Z\times\Omega}$, hence, by Proposition 3.3, the proof is finished. $\square$

By Proposition 3.10, for fiberwise well-pointed spaces $\mathbf{U}$ over Z and $\mathbf{U}'$ over Z' we can define

$$[\mathbf{U}]_Z \wedge [\mathbf{U}']_{Z'} := [\mathbf{U}\wedge\mathbf{U}']_{Z\times Z'}.$$

The role of the unit element for smash product is played by the homotopy type Σ^0 (also denoted $\overline{1}$), i.e. the homotopy type of punctured 0-dimensional sphere (S^0, s_0) over $\{s_0\}$. Up to obvious identifications, for every fiberwise well-pointed space $\mathbf{U}$ over Z,

$$[\mathbf{U}]_Z \wedge \Sigma^0 = \Sigma^0 \wedge [\mathbf{U}]_Z = [\mathbf{U}]_Z.$$

4 Fiberwise pointed spaces generated by closed pairs

Let X be a topological space and let $\omega : X \to Z$ be a continuous map.

For $P = (P_1, P_2)$ a pair of closed spaces, we define $U_\omega(P)$ as the adjunction $P_1 \cup_{\omega|_{P_2}} Z$ (see [E]), i.e.

$$U_\omega(P) := Z\times 0 \cup P_1 \times 1/\sim,$$

where $\sim$ denotes the equivalence relation given for $(u,i),(u',i') \in Z\times 0\cup P_1\times 1$ by

$$(u,i)\sim(u',i') \text{ if } i=i', u=u' \text{ or } i=1, i'=0, u\in P_2, u'=\omega(u).$$

The equivalence class of an element $(y,i)\in Z\times 0\cup P_1\times 1$ with respect to $\sim$ will be denoted by $[y,i]_{\omega,P}$ or briefly $[y,i]_P$. We define the quotient map by

$$q_{\omega,P} : Z\times 0\cup P_1\times 1 \ni (y,i) \to [y,i]_{\omega,P} \in U_\omega(P).$$

Proposition 4.1 *If $\omega|_{P_2}$ is a closed map then the quotient map $q_{\omega,P}$ is also closed.* □

Define $r_{\omega,P} : U_\omega(P) \to Z$ and $s_{\omega,P} : Z \to U_\omega(P)$ by

$$r_{\omega,P}([x,1]_{\omega,P}) := \omega(x),\ r_{\omega,P}([z,0]_{\omega,P}) := z$$
$$s_{\omega,P}(z) := [z,0]_{\omega,P}.$$

It is easy to verify that the maps $r_{\omega,P}, s_{\omega,P}$ are well defined and continuous and the triple $\mathbf{U}_\omega(P) := \mathbf{U}(P) := (U_\omega(P), r_{\omega,P}, s_{\omega,P})$ is a fiberwise pointed space over Z.

The following result is a straightforward consequence of definitions and facts presented at the end of Section 3.

Proposition 4.2 *If P is a cofiberd closed pair then $\mathbf{U}_\omega(P)$ is a fiberwise well-pointed space over Z.* □

If $P \subset Q$ are closed pairs in X then the inclusion induces an h.f.p. map (i.e. a morphism in Fib_Z):

$$\mathbf{U}_\omega(\iota_{PQ}) : \mathbf{U}_\omega(P) \longrightarrow \mathbf{U}_\omega(Q),\quad [u,i]_P \mapsto [u,i]_Q$$

Proposition 4.3 *If $P \subset Q$ are closed pairs such that $P_1 \setminus P_2 = Q_1 \setminus Q_2$ and $\omega|_{Q_2}$ is closed then $\mathbf{U}_\omega(\iota_{PQ})$ is an isomorphism in* Fib_Z.

Proof: In order to prove the result it suffices to show that $\mathbf{U}_\omega(\iota_{PQ})$ is a homeomorphism. Since $\mathbf{U}_\omega(\iota_{PQ})$ is a continuous bijective map, all what we need is to prove that $\mathbf{U}_\omega(\iota_{PQ}) \circ q_{\omega,P}$ is closed. Actually, the latter map is the composition of the inclusion $Z \times 0 \cup P_1 \times 1 \hookrightarrow Z \times 0 \cup Q_1 \times 1$ and $q_{\omega,Q}$. Since P_1 is closed, the inclusion is a closed map, and thus the assertion follows by Proposition 4.1. □

Now we present examples and results concerning weak homotopy type of the spaces introduced in this section. We begin with a trivial observation that for every map $\omega : X \to Z$,

$$[\mathbf{U}_\omega(\emptyset)]_Z = \overline{0}_Z.$$

More interesting weak homotopy types are described in the following:

Example 4.4 Let $C_1 = S^1$ and $C_2 = \{x : |x - (2,0)| = 1\}$ be two circles in the plane. Consider four closed pairs in $\mathrm{R}^2 \setminus 0$:

$$P_1 = (C_1, \emptyset),\ P_2 = (C_2, \emptyset),\ P_3 = (C_1, \{(1,0)\}),\ P_4 = (C_2, \{(3,0)\}).$$

Putting $\omega = \mathrm{id}_{\mathrm{R}^2\setminus 0}$, we get $[\mathbf{U}(P_i)]_{\mathrm{R}^2\setminus 0} \neq [\mathbf{U}(P_3)]_{\mathrm{R}^2\setminus 0}$ for $i = 1, 2$, because P_3 is connected. By Proposition 3.5, we also have $[\mathbf{U}(P_1)]_{\mathrm{R}^2\setminus 0} \neq [\mathbf{U}(P_2)]_{\mathrm{R}^2\setminus 0}$. However, one can show that $[\mathbf{U}(P_3)]_{\mathrm{R}^2\setminus 0} = [\mathbf{U}(P_4)]_{\mathrm{R}^2\setminus 0}$.

The following example shows that for a given pair P the weak homotopy types of $\mathbf{U}_\omega(P)$ can be distinct for various maps ω from the same homotopy class.

Example 4.5 Let $X = Z = \mathrm{R}$ and let $P = (P_1, P_2)$ be given by

$$P_1 = \{\frac{1}{n} : n \in \mathrm{Z}^+\} \cup \{0\},\quad P_2 = \{0\}.$$

Denote by τ_1 the translation $x \mapsto x + 1$. Then $\tau_1 \simeq \mathrm{id}_{\mathrm{R}}$, but

$$[\mathbf{U}_{\tau_1}(P)]_{\mathrm{R}} \neq [\mathbf{U}_{\mathrm{id}_{\mathrm{R}}}(P)]_{\mathrm{R}}.$$

This can be shown by essentially the same argument as in Example 3.8.

In the next proposition we indicate that $[\mathbf{U}_\omega(P)]_Z$ depends only on the homotopy class of ω if an additional assumption on the pair P is imposed. Actually, we prove an even stronger result. Let $\phi, \psi : X \to Z$ be continuous maps.

Proposition 4.6 *If $P = (P_1, P_2)$ is a cofiberd closed pair and $\phi|_{P_2} \simeq \psi|_{P_2} : P_2 \to Z$ then*

$$[\mathbf{U}_\phi(P)]_Z = [\mathbf{U}_\psi(P)]_Z.$$

Proof: Let (G, a) be a Strøm structure for P and let $\omega_t : P_2 \to Z$, $t \in I$, be a homotopy joining $\omega_0 = \phi|_{P_2}$ with $\omega_1 = \psi|_{P_2}$. Define maps $f : \mathbf{U}_{\omega_0}(P) \to \mathbf{U}_{\omega_1}(P)$ and $f' : \mathbf{U}_{\omega_1}(P) \to \mathbf{U}_{\omega_0}(P)$ by

$$f([z, 0]_{\omega_0, P}) := [z, 0]_{\omega_1, P},$$

$$f([x,1]_{\omega_0,P}) := \begin{cases} [\omega_{2a(x)}G_1(x),0]_{\omega_1,P} & \text{if } x \in \operatorname{cl} a^{-1}([0,\frac{1}{2})), \\ [G_{2-2a(x)}(x),1]_{\omega_1,P} & \text{if } \frac{1}{2} \le a(x), \end{cases}$$

$$f'([z,0]_{\omega_1,P}) := [z,0]_{\omega_0,P},$$

$$f'([x,1]_{\omega_1,P}) := \begin{cases} [\omega_{1-2a(x)}G_1(x),0]_{\omega_0,P} & \text{if } x \in \operatorname{cl} a^{-1}([0,\frac{1}{2})), \\ [G_{2-2a(x)}(x),1]_{\omega_0,P} & \text{if } \frac{1}{2} \le a(x). \end{cases}$$

Both the maps f and f' are h.f.p. as one can directly check. Define a homotopy $h : U_{\omega_0}(P) \times I \to U_{\omega_0}(P)$ by

$$h([z,0]_{\omega_0,P},t) := [z,0]_{\omega_0,P},$$

$$h([x,1]_{\omega_0,P},t) := \begin{cases} [\omega_{2ta(x)}G_1(x),0]_{\omega_0,P} & \text{if } x \in \operatorname{cl} a^{-1}([0,\frac{1}{2})), \\ [\omega_{t(1-2a(G_{2-2a(x)}(x))}G_1G_{2-2a(x)}(x),0]_{\omega_0,P} & \text{if } \frac{1}{2} \le a(x), \\ & G_{2-2a(x)}(x) \in \operatorname{cl} a^{-1}([0,\frac{1}{2})), \\ [G_{2-2a(G_{2-2a(x)}(x))}G_{2-2a(x)}(x),1]_{\omega_0,P} & \text{if } \frac{1}{2} \le a(x), \\ & \frac{1}{2} \le aG_{2-2a(x)}(x). \end{cases}$$

Thus $f' \circ f \simeq h_0 \operatorname{rel} s_{\omega_0,P}(Z)$. Define another homotopy $j : U_{\omega_0}(P) \times I \to U_{\omega_0}(P)$ by

$$j([z,0]_{\omega_0,P},t) := [z,0]_{\omega_0,P},$$

$$j([x,1]_{\omega_0,P},t) := \begin{cases} [\omega_0G_1(x),0]_{\omega_0,P} & \text{if } x \in \operatorname{cl} a^{-1}([0,\frac{1}{2})), \\ [\omega_0G_1G_{2-2a(x)}(x),0]_{\omega_0,P} & \text{if } \frac{1}{2} \le a(x), \\ & G_{2-2a(x)}(x) \in \operatorname{cl} a^{-1}([0,\frac{1}{2})), \\ [G_{t+(1-t)(2-2a(G_{2-2a(x)}(x)))}G_{2-2a(x)}(x),1]_{\omega_0,P} & \text{if } \frac{1}{2} \le a(x), \\ & \frac{1}{2} \le aG_{2-2a(x)}(x), \end{cases}$$

hence $h_0 \simeq j_1 \operatorname{rel} s_{\omega_0,P}(Z)$. The next homotopy $k : U_{\omega_0}(P) \times I \to U_{\omega_0}(P)$ is given by

$$k([z,0]_{\omega_0,P},t) := [z,0]_{\omega_0,P},$$

$k([x,1]_{\omega_0,P},t) :=$

$$\begin{cases} [\omega_0 G_1 G_t(x),0]_{\omega_0,P} & \text{if } a(x) \le \frac{1}{2}, \\ [\omega_0 G_1 G_{t(2-2a(x))}(x),0]_{\omega_0,P} & \text{if } \frac{1}{2} \le a(x), aG_{t(2-2a(x))}(x) \le \frac{1}{2}, \\ [G_1 G_{t(2-2a(x))}(x),1]_{\omega_0,P} & \text{if } \frac{1}{2} \le a(x), \frac{1}{2} \le aG_{2-2a(x)}(x). \end{cases}$$

Since G comes from the Strøm structure, $G_1 G_1(x) = G_1(x)$ if $a(x) < 1$, hence $k_1 = j_1$ and thus $j_1 \simeq k_0 \operatorname{rel} s_{\omega_0,P}(Z)$. The last homotopy $m : U_{\omega_0}(P) \times I \to U_{\omega_0}(P)$ is simply defined by $m_t([z,0]_{\omega_0,P}) = [z,0]_{\omega_0,P}$, $m_t([x,1]_{\omega_0,P}) = [G_t(x),1]_{\omega_0,P}$. It is easy to check that $k_0 = m_1$, hence $k_0 \simeq \mathrm{id}_{U_{\omega_0}(P)} \operatorname{rel} s_{\omega_0,P}(Z)$. By composing the above homotopies we conclude that $f' \circ f \simeq \mathrm{id}_{U_{\omega_1}(P)} \operatorname{rel} s_{\omega_0,P}(Z)$. Similarly, one can prove that also $f \circ f'$ is homotopic to the identity, hence, by Proposition 3.3, the proof is finished. □

Let z_0 be a point in Z and let $c := c_{z_0} : X \to Z$, $x \mapsto z_0$, be a constant map. If P is a closed pair then $U_c(P) = P_1/P_2 \vee (Z, z_0)$ and $r_{c,P}([x,1]_{c,P}) = z_0$ for every $x \in P_1$. In that case the weak homotopy type $[\mathbf{U}_c(P)]_Z$ is called *simple*.

Corollary 4.7 *If $P = (P_1, P_2)$ is cofiberd and P_1 is contractible in X then $[\mathbf{U}_\omega(P)]_Z$ is simple for every $\omega : X \to Z$.* □

Let P be a closed pair in X and let $\zeta : Z \to Z'$ be a continuous map. Define $\zeta_P : U_\omega(P) \to U_{\zeta\circ\omega}(P)$ by

$$\zeta_P([z,0]_{\omega,P}) = [\zeta(z),0]_{\zeta\circ\omega,P}, \quad \zeta_P([x,1]_{\omega,P}) = [x,1]_{\zeta\circ\omega,P}.$$

Since $\zeta_P \circ q_{\omega,P} = q_{\zeta\circ\omega,P} \circ (\zeta \times 0 \cup \mathrm{id}_{P_1\times 1})$, the map ζ_P is continuous. Assume that Q is another closed pair and let $f : \mathbf{U}_\omega(P) \to \mathbf{U}_\omega(Q)$ be an h.f.p. map. Define $f' : U_{\zeta\circ\omega}(P) \to U_{\zeta\circ\omega}(Q)$ by

$$f'([z',0]_{\zeta\circ\omega,P}) = [z',0]_{\zeta\circ\omega,Q}, \quad f'([x,1]_{\zeta\circ\omega,P}) = (\zeta_Q \circ f)([x,1]_{\omega,P}).$$

The correctnes of the definition of f' is a consequence of the fact that $f([z,0]_{\omega,P}) = [z,0]_{\omega,Q}$ for every $z \in Z$ (recall that f is h.f.p.). The map f' is continuous because the restrictions $(f' \circ q_{\zeta\circ\omega,P})|_{Z'\times 0}$ and $(f' \circ q_{\zeta\circ\omega,P})|_{P_1\times 1}$ are continuous, which follows from the description of f'. Moreover, f' is also h.f.p. Indeed, let $h : U_\omega(P) \times I \to Z$ be a homotopy relative to $s_{\omega,P}(Z)$ joining $r_{\omega,Q} \circ f$ with $r_{\omega,P}$. Define $h' : U_{\zeta\circ\omega}(P) \times I \to Z'$ by

$$h'([z',0]_{\zeta\circ\omega,P},t) = z', \quad h'([x,1]_{\zeta\circ\omega,P},t) = (\zeta \circ h)([x,1]_{\omega,P},t).$$

It follows by Remark 3.9 and an argument similar to the above that h' is correctly defined and continuous, and is the required homoptopy joining $r_{\zeta\circ\omega,Q}\circ f$ with $r_{\zeta\circ\omega,P}$, hence $f' : \mathbf{U}_{\zeta\circ\omega}(P) \to \mathbf{U}_{\zeta\circ\omega}(Q)$ is h.f.p. By a suitable modification of the above arguments one can show that if $f \simeq g : \mathbf{U}_\omega(P) \to \mathbf{U}_\omega(Q)$ in Fib_Z then also the corresponding h.f.p. maps $f', g' : \mathbf{U}_{\zeta\circ\omega}(P) \to \mathbf{U}_{\zeta\circ\omega}(Q)$ are homotopic in $\mathrm{Fib}_{Z'}$. As a conclusion we get the following:

Proposition 4.8 *If* $[\mathbf{U}_\omega(P)]_Z = [\mathbf{U}_\omega(Q)]_Z$ *and* $\zeta : Z \to Z'$ *is continuous, then* $[\mathbf{U}_{\zeta\circ\omega}(P)]_{Z'} = [\mathbf{U}_{\zeta\circ\omega}(Q)]_{Z'}$. □

As immediate corollaries from the above proposition we get results which assert that the information provided by $[\mathbf{U}_\omega(\cdot)]_Z$ is maximal in the case $\omega = \mathrm{id}_X$ and minimal if ω is a constant map onto a one-point space.

Corollary 4.9 *If* $[\mathbf{U}_\omega(P)]_Z = [\mathbf{U}_\omega(Q)]_Z$ *for some* $\omega : X \to Z$ *then* $[P_1/P_2] = [Q_1/Q_2]$. □

Corollary 4.10 *If* $[\mathbf{U}_{\mathrm{id}_X}(P)]_X = [\mathbf{U}_{\mathrm{id}_X}(Q)]_X$ *then for every space* Z *and every continuous map* $\omega : X \to Z$, $[\mathbf{U}_\omega(P)]_Z = [\mathbf{U}_\omega(Q)]_Z$. □

Propositions 4.6 and 4.8 imply:

Corollary 4.11 *If* P *and* Q *are cofiberd pairs, and* $\zeta : Z \to Z'$ *is a homotopy equivalence then* $[\mathbf{U}_\omega(P)]_Z = [\mathbf{U}_\omega(Q)]_Z$ *if and only if* $[\mathbf{U}_{\zeta\circ\omega}(P)]_{Z'} = [\mathbf{U}_{\zeta\circ\omega}(Q)]_{Z'}$. □

We conclude this section by results on sums and products of closed pairs. Let $P = (P_1, P_2)$ and $Q = (Q_1, Q_2)$ be disjoint closed pairs in X, i.e. $P_1 \cap Q_1 = \emptyset$. Then

$$\mathbf{U}_\omega(P \cup Q) = \mathbf{U}_\omega(P) \vee \mathbf{U}_\omega(Q),$$

hence:

Proposition 4.12 *If* P *and* Q *are disjoint closed pairs in* X *then*

$$[\mathbf{U}_\omega(P \cup Q)]_Z = [\mathbf{U}_\omega(P)]_Z \vee [\mathbf{U}_\omega(Q)]_Z.\square$$

Now let P be a closed pair in X and P' in X', and let $\omega : X \to Z$ and $\omega' : X' \to Z'$ be continuous. Recall that $P \times P' = (P_1 \times P_1', P_1 \times P_2' \cup P_2 \times P_1')$. Then, up to a natural isomorphism,

$$\mathbf{U}_{\omega\times\omega'}(P \times P') = \mathbf{U}_{\omega}(P) \wedge \mathbf{U}_{\omega'}(P').$$

As a consequence of Proposition 4.2 we get:

Proposition 4.13 *If P and P' are cofiberd closed pairs then*

$$[\mathbf{U}_{\omega\times\omega'}(P \times P')]_{Z\times Z'} = [\mathbf{U}_{\omega}(P)]_Z \wedge [\mathbf{U}_{\omega'}(P')]_{Z'}.\square$$

5 Regular index pairs and isolating blocks.

Let X be a metrizable locally compact space and let the map $\varphi : X \times \mathrm{R}^+ \ni (x,t) \to x \cdot t \in X$ be a semiflow. We recall several concepts from the theory of isolated invariant sets, cf. [Co], [Ch], [Ry], and [Sm].

Let I be an interval in R with $0 \in I$ and let $A \subset X$. A single valued mapping $\sigma : I \to A$ is called a *solution for φ through $x \in A$* if $\sigma(t+s) = \varphi(\sigma(t), s)$ for all $t, t+s \in I, s > 0$, and $\sigma(0) = x$.

Given an arbitrary set $A \subset X$, we define the *maximal invariant part of A* by

$$\operatorname{Inv} A := \{x \in A : \exists \sigma : \mathrm{R} \to A \text{ a solution for } \varphi \text{ through } x\}$$

the *exit function on A* by

$$\sigma_A : A \ni x \longrightarrow \sup\{t \geq 0 \mid x \cdot [0,t] \subset A\} \in [0, \infty],$$

and the *exit set* $A^- := \sigma_A^{-1}(0)$.

A set $S \subset X$ is called an *isolated invariant set* if S is compact and there exists U, a neighborhood of S, such that $S = \operatorname{Inv} U$. A compact subset $N \subset X$ is called an *isolating neighborhood for φ* if $\operatorname{Inv} N \subset \operatorname{int} N$. It is easy to check that if N is an isolating neighborhood then $\operatorname{Inv} N$ is an isolated invariant set. An isolating neighborhood B is called an *isolating block* iff B^- is closed.

Proposition 5.1 (Ważewski) *An isolating neighborhood B is an isolating block if and only if the map σ_B is continuous.*

Let N be an isolating neighborhood for φ. A pair $P = (P_1, P_2)$ of compact subsets $P_2 \subset P_1 \subset N$ is called an *index pair in* N if the following conditions are satisfied:

$$x \in P_i, t > 0, x \cdot [0,t] \subset N \;\Rightarrow\; x \cdot [0,t] \subset P_i \tag{5.1}$$

$$x \in P_1, t > 0, x \cdot t \notin N \;\Rightarrow\; \exists t' \in [0,t]\; x \cdot t' \in P_2, x \cdot [0,t'] \subset N \tag{5.2}$$

$$\operatorname{Inv} N \subset \operatorname{int}(P_1 \setminus P_2) \tag{5.3}$$

For an index pair P we define functions $\sigma_P, \tau_P : P_1 \to [0,\infty]$ by

$$\sigma_P(x) := \begin{cases} \sigma_{\operatorname{cl}(P_1 \setminus P_2)}(x) & \text{if } x \in \operatorname{cl}(P_1 \setminus P_2) \\ 0 & \text{if } x \in \operatorname{int}_{P_1} P_2 \end{cases} \tag{5.4}$$

$$\tau_P(x) := \begin{cases} \sup\{t \geq 0 \mid x \cdot [0,t] \subset P_1 \setminus P_2\} & \text{if } x \in P_1 \setminus P_2 \\ 0 & \text{if } x \in P_2 \end{cases} \tag{5.5}$$

Proposition 5.2 *σ_P is upper semicontinuous, τ_P is lower semicontinuous and $\tau_P(x) \leq \sigma_P(x)$ for $x \in P_1$.*

One easily verifies the following proposition.

Proposition 5.3 *If φ is a flow, then the following three conditions are equivalent.*

1. *σ_P is continuous.*
2. *τ_P is continuous.*
3. *$\sigma_P = \tau_P$.*

This proposition is not true for semiflows as the example in Figure 5 illustrates. The large rectangle is P_1, and the shaded area is P_2.

We call the index pair P *regular* if $\tau_P(x) = \sigma_P(x)$ for $x \in P_1$. As an immediate consequence of Proposition 5.2 we get

Proposition 5.4 *If P is a regular index pair, then $\sigma_P = \tau_P$ is continuous and $\operatorname{cl}(P_1 \setminus P_2)$ is an isolating block.*



Figure 2: An index pair for a semiflow with $\sigma_P \neq \tau_P$.

A convenient way of constructing regular index pairs or isolating blocks is by means of Lyapunov functions. A function $\alpha : A \to [0, \infty]$ is called *strictly increasing (decreasing) along the trajectories of* φ if for any $x \in A$ and $t > 0$ such that $x \cdot [0, t] \subset A$ and $\alpha(x) > 0$ there is $\alpha(x \cdot t) > \alpha(x)$ $(\alpha(x \cdot t) < \alpha(x))$. A pair of functions $\mu = (\mu_1, \mu_2)$, where $\mu_i : N \to [0, \infty]$ for $i = 1, 2$ will be called a *Lyapunov pair* in N if μ_1 is strictly decreasing, μ_2 is strictly increasing along the trajectories of φ and $\mathrm{Inv}^N = \mu_1^{-1}(0) \cap \mu_2^{-1}(0)$. The following theorem is a by-product of the proof of Theorem 1.5.1 in [Ry].

Theorem 5.5 *If N is an isolating neighborhood for S which is sufficiently close to S then it admits a Lyapunov pair.*

A standard compactness argument leads to the following

Proposition 5.6 *Assume $U \subset N$ is an open neighborhood of $S = \mathrm{Inv}\, N$ and μ is a Lyapunov pair in N. Then there exists an $\epsilon > 0$ such that*

$$\mu_1^{-1}([0, \epsilon]) \cap \mu_2^{-1}([0, \epsilon]) \subset U. \tag{5.6}$$

Lyapunov pairs may be used to construct regular index pairs. Let N be an isolating neighborhood which admits a Lyapunov pair μ. Let $U \subset \mathrm{int}\, N$ be open and let $\epsilon > 0$ be such that (5.6) is satisfied. Put $\alpha_1, \alpha_2 \in (0, \epsilon)$ and for $\alpha := (\alpha_1, \alpha_2)$ define

$$P^\alpha := (P_1^\alpha, P_2^\alpha),$$

where

$$P_1^\alpha := \{\, x \in N \mid \mu_1(x) \le \alpha_1 \,\} \tag{5.7}$$

$$P_2^\alpha := \{\, x \in P_1^\alpha \mid \mu_2(x) \ge \alpha_2 \,\}. \tag{5.8}$$

We have the following

Theorem 5.7 *P^α is a regular index pair in N which satisfies $P_1^\alpha \backslash P_2^\alpha \subset U$.*

Proof: Property (5.1) is obvious. To see (5.2) take $x \in P_1^\alpha$ and $t > 0$ such that $x \cdot t \notin N$. Put $t' := \sigma_N(x)$. Then $x \cdot t' \in \operatorname{bd} N$, i.e. $\mu_2(x \cdot t') \ge \epsilon \ge \alpha_2$. This proves (5.2). Property (5.3) is obvious. It remains to show that P^α is regular. We have

$$P_1^\alpha \setminus P_2^\alpha = \{\, x \in N \mid \mu_1(x) \le \alpha_1, \mu_2(x) < \alpha_2 \,\}$$

and

$$\operatorname{cl}(P_1^\alpha \setminus P_2^\alpha) \subset \{\, x \in N \mid \mu_1(x) \le \alpha_1, \mu_2(x) \le \alpha_2 \,\}.$$

Thus if $x \in P_1$ we get $\mu_2(x \cdot \tau_P(x)) = \alpha_2$ and $\mu_2(x \cdot \sigma_P(x)) \le \alpha_2$. Since μ_2 is increasing along solutions, it must be $\sigma_P(x) \le \tau_P(x)$. The other inequality follows from Proposition 5.2. □

As a consequence we obtain the following proposition.

Lemma 5.8 *Assume N is an isolating neighborhood which admits a Lyapunov pair. Then for any $n \in N$ there exists a sequence of regular index pairs $P^1, P^2, \ldots P^n$ in N such that $P^{i-1} \subset \operatorname{int} P^i$ for $i = 2, 3, \ldots n$.*

Proof: Choose numbers $0 < \alpha_1^1 < \alpha_1^2 < \ldots \alpha_1^n \epsilon$, $\epsilon > \alpha_1^2 > \alpha_2^2 > \ldots \alpha_2^n > 0$ and put $P^i := P^{\alpha^i}$, where $\alpha^i := (\alpha_1^i, \alpha_2^i)$. It now straightforward to verify that the assertion is satisfied. □

Proposition 5.9

(i) if P and Q are regular index pairs in N, then so is $P \cap Q := (P_1 \cap Q_1, P_2 \cap Q_2)$.

(ii) If $P \subseteq Q$ are regular index pairs for N, then so are $Q' := (P_1, P_1 \cap Q_2)$ and $Q" := (P_1 \cup Q_2, Q_2)$.

Proof: It is an easy exercise to show that $P \cap Q, Q', Q"$ satisfy properties ((5.1)-(5.3)). To show regularity of $P \cap Q, Q', Q"$ observe first that it follows easily from (5.5) that

$$P \subset Q \;\Rightarrow\; \tau_P \geq \tau_{Q|P_1}. \tag{5.9}$$

We will show that

$$\tau_{P\cap Q} = \max(\tau_{P|P_1\cap Q_1}, \tau_{Q|P_1\cap Q_1}).$$

One inequality follows immediately from 5.9. Assume the other is not true. Then there exists an $x \in P_1 \cap Q_1$ and a $t \in \mathrm{R}^+$ such that

$$\tau_{P\cap Q}(x) > t > \max(\tau_P(x), \tau_Q(x))$$

In particular we get from (5.5) that

$$x \cdot [0,t] \cap P_2 \cap Q_2 = \emptyset$$

and for some $t_1, t_2 \in [0,t]$

$$x \cdot t_1 \in P_2,\; x \cdot t_2 \in Q_2$$

It follows from (5.1) that $x \cdot \max(t_1, t_2) \in P_2 \cap Q_2$, a contradiction. Thus $\tau_{P\cap Q}$ is continuous as a maximum of two continuous functions and consequently $P \cap Q$ is a regular index pair.

A similar argument shows that $\tau_{Q'} = \tau_{Q|P_1}$ and $\tau_{Q"} = \tau_{Q|P_1\cup Q_2}$ and the conclusion follows. □

6 The Conley index over a base

In the reminder of this paper we assume that $\omega : X \to Z$ is continuous and Z is a Hausdorff space. We connect results on flows from the previous section with the topological background from Section 4. First observe that a regular index pair is cofiberd. This is an immediate consequence of 5.4 and [Ry, Th. 3.7], Actually, it can be easily verified by constructing a Strøm structure using the exit function; similar constructions will frequently appear in the reminder of this paper. As a consequence of Proposition 4.2, we have:

Proposition 6.1 *If P is a regular index pair then $\mathbf{U}_\omega(P)$ is a fiberwise well-pointed space over Z.* $\square$

We will prove the following two theorems. The second clearly implies Theorem 2.1.

Theorem 6.2 *Let N be an isolating neighborhood and let $P \subset Q$ be two regular index pairs in N. Then $\mathbf{U}_\omega(\iota_{PQ})$ is a weak fiberwise homotopy equivalence.*

Theorem 6.3 *Let S be an isolated invariant set. Then $h_\omega(S,\varphi) := [\mathbf{U}_\omega(P)]_Z$ is independent of the choice of an isolating neighborhood N of S and of a regular index pair P in N.*

The homotopy type $h_\omega(S,\varphi)$ given by the above theorem (denoted shortly by $h_\omega(S)$ if φ is clear from the context) will be called the *Conley index of S over base Z with base map ω* or just the *Conley index of S over Z* if ω is clear from context.

Proof of Theorem 6.2 The proof proceeds in two steps.

Step 1 We make an extra assumption that P and Q differ by at most one coordinate, i.e. $P_1 = Q_1$ or $P_2 = Q_2$. Let us assume first that $P_1 = Q_1$. Fix an $\epsilon > 0$. For $x \in Q_1$ put $\alpha(x) := \sigma_P(x) - \sigma_Q(x)$ and define the map $f : \mathbf{U}_\omega(Q) \to \mathbf{U}_\omega(P)$ by

$$f([x,i]_Q) := \begin{cases} [\omega\Big(x \cdot (\sigma_Q(x) + \frac{\sigma_Q(x)}{\epsilon}\alpha(x)\Big), 0]_P & \text{if } i = 1,\ 0 \le \sigma_Q(x) \le \epsilon \\ [x \cdot \frac{2\epsilon - \sigma_Q(x)}{\epsilon}(\epsilon + \alpha(x)), 1]_P & \text{if } i = 1,\ \epsilon \le \sigma_Q(x) \le 2\epsilon \\ [x,1]_P & \text{if } i = 1,\ \sigma_Q(x) \ge 2\epsilon \\ [x,0]_P & \text{if } i = 0 \end{cases}$$

It is a lengthy but straightforward task to verify that the map is well defined and continuous. We will show that f is a homotopy inverse of $\mathbf{U}_\omega(\iota_{PQ})$. For this end consider the homotopy $F_\lambda : \mathbf{U}_\omega(Q) \to \mathbf{U}_\omega(Q)$ given by

$$F_\lambda([x,i]_Q) := \begin{cases} [\omega\Big(x \cdot (\sigma_Q(x) + \frac{\sigma_Q(x)}{\epsilon}\alpha(x)\Big), 0]_Q & \text{if } i = 1,\ 0 \le \sigma_Q(x) \le \lambda\epsilon \\ [x \cdot \frac{2\epsilon - \sigma_Q(x)}{\epsilon}(\epsilon + \alpha(x))\frac{\lambda}{2-\lambda}, 1]_Q & \text{if } i = 1,\ \lambda\epsilon \le \sigma_Q(x) \le 2\epsilon \\ [x,1]_Q & \text{if } i = 1,\ \sigma_Q(x) \ge 2\epsilon \\ [x,0]_Q & \text{if } i = 0 \end{cases}$$

Again, the map is well defined and continuous. Moreover, one easily verifies that $F_0 = \mathrm{id}_{\mathbf{U}_\omega(Q)}$ and $F_1 = \mathbf{U}_\omega(\iota_{PQ})f$, which means that f is the right homotopy inverse of $\mathbf{U}_\omega(\iota_{PQ})$. To show that it is also the left inverse first observe that the map f also satisfies the formula

$$f([x,i]_Q) := \begin{cases} [\omega(x),0]_P & \text{if } i=1,\ 0 \le \sigma_P(x) \le \alpha(x) \\ [\omega\big(x \cdot (\sigma_P(x) + \frac{\sigma_Q(x)}{\epsilon}\alpha(x)\big),0]_P & \text{if } i=1,\ \alpha(x) \le \sigma_P(x) \le \epsilon + \alpha(x) \\ [x \cdot \frac{2\epsilon - \sigma_Q(x)}{\epsilon}(\epsilon+\alpha(x)),1]_P & \text{if } i=1,\ \epsilon+\alpha(x) \le \sigma_P(x) \le 2\epsilon+\alpha(x) \\ [x,1]_P & \text{if } i=1,\ \sigma_P(x) \ge 2\epsilon+\alpha(x) \\ [x,0]_P & \text{if } i=0 \end{cases}$$

and consider the homotopy $G_\lambda : \mathbf{U}_\omega(P) \to \mathbf{U}_\omega(P)$ given by

$$G_\lambda([x,i]_Q) := \begin{cases} [\omega(x),0]_P & \text{if } i=1,\ 0 \le \sigma_P(x) \le \lambda\alpha(x) \\ [\omega\big(x \cdot (\frac{\epsilon+\alpha(x)}{\epsilon}(\sigma_P(x) - \lambda\alpha(x))\big),0]_P & \text{if } i=1,\ \lambda\alpha(x) \le \sigma_P(x) \le \lambda(\epsilon+\alpha(x)) \\ [x \cdot \frac{2\epsilon+\alpha(x)-\sigma_P(x)}{2\epsilon+\alpha(x)+\lambda(\epsilon+\alpha(x)}\lambda(\epsilon+\alpha(x)),1]_P & \text{if } i=1,\ \lambda(\epsilon+\alpha(x)) \le \sigma_P(x) \le 2\epsilon+\alpha(x) \\ [x,1]_P & \text{if } i=1,\ \sigma_P(x) \ge 2\epsilon+\alpha(x) \\ [x,0]_P & \text{if } i=0 \end{cases}$$

One can verify that the map is well defined and continuous and $F_0 = \mathrm{id}_{\mathbf{U}_\omega(P)}$ and $F_1 = f\mathbf{U}_\omega(\iota_{PQ})$. This means that f is a homotopy inverse of $\mathbf{U}_\omega(\iota_{PQ})$. It follows from Proposition 3.3 that f is a representative of the inverse of $[\mathbf{U}_\omega(\iota_{PQ})]_Z$ in HtpFib_Z, which proves the case when $P_1 = Q_1$.

Assume in turn that $P_2 = Q_2$. There exists a $T > 0$ such that for every $x \in Q_1$ we have $x \cdot [0,T] \neq \emptyset$. Put $\beta(x) := \min(T, \sigma_Q(x))$ and define the function $g : \mathbf{U}_\omega(Q) \to \mathbf{U}_\omega(P)$ by

$$g([u,i]_Q) := \begin{cases} [u \cdot \beta(u),1]_P & \text{if } i=1 \\ [u,0]_P & \text{if } i=0 \end{cases}$$

One can easily verify that g is a well defined continuous function. We will show that g is a weak homotopy inverse of $\mathbf{U}_\omega(\iota_{PQ})$. For this end consider the homotopy $F_\lambda : \mathbf{U}_\omega(Q) \to \mathbf{U}_\omega(Q)$ given by

$$F_\lambda([u,i]_Q) := \begin{cases} [u \cdot \lambda\beta(u),1]_Q & \text{if } i=1 \\ [u,0]_Q & \text{if } i=0 \end{cases}$$

and the homotopy $G_\lambda : \mathbf{U}_\omega(P) \to \mathbf{U}_\omega(P)$ given by

$$G_\lambda([u,i]_P) := \begin{cases} [u \cdot \lambda\beta(u),1]_P & \text{if } i=1 \\ [u,0]_P & \text{if } i=0 \end{cases}$$

One easily checks that both homotopies are well defined and join respectively $\mathrm{id}_{\mathbf{U}_\omega(Q)}$ with $\mathbf{U}_\omega(\iota_{PQ})g$ and $\mathrm{id}_{\mathbf{U}_\omega(P)}$ with $g\mathbf{U}_\omega(\iota_{PQ})$. Thus Proposition 3.3 shows that $\mathbf{U}_\omega(P)$ and $\mathbf{U}_\omega(Q)$ are weakly fiberwise homotopic. This completes the proof of Step 1.

Step 2 We lift the extra assumption. Put $R_1 := P_1 \cup Q_2, R_2 := P_1 \cap Q_2$. By Proposition 5.9 (ii),(P_1, R_2) and (R_1, Q_2) are index pairs. We have the commutative diagram of inclusions

$$\begin{array}{ccc} (P_1, R_2) & \xrightarrow{j_2} & (R_1, Q_2) \\ \uparrow{\scriptstyle i_1} & & \downarrow{\scriptstyle i_3} \\ (P_1, P_2) & \xrightarrow{j} & (Q_1, Q_2) \end{array}$$

It is clear that the pair of index pairs $(P_1, P_2) \subset (P_1, R_2)$ satisfies the extra assumption of Step 1 and so does $(R_1, Q_2) \subset (Q_1, Q_2)$. Thus the inclusions i_1 and i_3 induce isomorphisms in HtpFib_Z . Since $P_1 \backslash R_2 = P_1 \backslash Q_2 = R_1 \backslash Q_2$, the inclusion j_2 induces an isomorphism by Proposition 4.3 . □

Proof of Theorem 6.3 We need to show that if M and N are two isolating neighborhoods of S, P a regular index pair for N and Q a regular index pair for M then $[\mathbf{U}_\omega(P)]_Z$ and $[\mathbf{U}_\omega(Q)]_Z$ have the same weak fiberwise homotopy type.

If $M = N$, then, y Proposition 5.9 (i), $P \cap Q$ is a regular index pair in N. Thus Theorem 6.2 may be applied to pairs $P \cap Q \subset P$ and $P \cap Q \subset Q$.

If $M \neq N$, one may always assume that $M \subset N$ since otherwise $M \cap N$ can be considered which is also an isolating neighborhood of S. Thus it is sufficient to show the existence of one regular index pair P in N and one regular index pair Q in M such that $[\mathbf{U}_\omega(P)]_Z$ and $[\mathbf{U}_\omega(Q)]_Z$ are isomorphic.

By Theorem 5.7 there exists a regular index pair P in N such that $P_1 \backslash P_2 \subset \mathrm{int}\, M$. It is easily verified that $Q := (M \cap P_1, M \cap P_2)$ is a regular index pair for M. Since $Q_1 \backslash Q_2 = M \cap (P_1 \backslash P_2) = P_1 \backslash P_2$, the inclusion $Q \subseteq P$ induces an isomorphism by Proposition 4.3 □

7 The properties of the Conley index over a base

As in the previous section, here we assume that S is an isolated invariant set for the semiflow φ on X. Recall that $\omega : X \to Z$ is continuous, and Z is

Hausdorff.

Proposition 7.1 *For any flow φ,*

$$h_\omega(\emptyset, \varphi) = \overline{0}_Z.$$

Proof: Use the regular index pair $(\emptyset, \emptyset)$. □

Proposition 7.2 *Suppose S and S' are disjoint isolated invariant sets. Then*

$$h_\omega(S \cup S', \varphi) = h_\omega(S, \varphi) \vee h_\omega(S', \varphi).$$

Proof: One can choose disjoint isolating neighborhoods for S and S', so the union is isolated. Now choose disjoint regular index pairs P for S and P' for S'. The result follows from Proposition 4.12. □

Let φ be as above and let φ' be a semiflow on X'. By $\varphi^\#$ denote the product flow on $X \times X'$, i.e. $\varphi_t^\#(x, x') = (\varphi_t(x), \varphi_t'(x'))$. Let $\omega : X \to Z$ and $\omega' : X' \to Z'$ be continuous functions.

Proposition 7.3 *Assume that S and S' are isolated invariant sets; S for φ and S' for φ'. Then*

$$h_{\omega \times \omega'}(S \times S', \varphi^\#) = h_\omega(S, \varphi) \wedge h_{\omega'}(S', \varphi').$$

Proof: This is a consequence of Propositions 4.13 and 6.1. □

Let $\varphi^\lambda : X \times \mathrm{R}^+ \to X$ be a a family of semiflows depending continuously on $\lambda \in [0, 1]$. We now prove the continuation theorem.

Proof of Theorem 2.2 Obviously it is sufficient to show that for any $\lambda_0 \in \Lambda$ there exists a neighborhood $\Delta_0 \subset \Lambda$ of λ_0 such that $h_\omega(N, \varphi^\lambda)$ does not depend on $\lambda \in \Delta_0$. Thus let us fix a $\lambda_0 \in \Lambda$. We introduce first some notation. If $\Delta \subset [0, 1]$ and $A \subset X \times I$ then we will write A^Δ for the set $A \cap X \times \Delta$ and we extend this notation in the obvious way to pairs, i.e. if $P = (P_1, P_2)$ then $P^\Delta := (P_1^\Delta, P_2^\Delta)$.

For $\Delta \subset [0, 1]$ there is a semiflow φ^Δ on $X \times \Delta$ given by

$$\varphi^\Delta : X \times \Delta \times \mathrm{R}^+ \ni ((x, \lambda), t) \to (\varphi^\lambda(x, t), \lambda) \in X \times \Delta.$$

Since

$$\mathrm{Inv}(N \times \Delta, \varphi^{\Delta}) = \bigcup_{\lambda \in \Delta} \mathrm{Inv}(N, \varphi^{\lambda}) \times \{ \lambda \} \subset \bigcup_{\lambda \in \Delta} \operatorname{int} N \times \{ \lambda \} = \operatorname{int}_{X \times \Delta} N \times \Delta,$$

we see that $N \times \Delta$ is an isolating neighborhood with respect to φ^{Δ}.

Let $M \subset N \times \Lambda$ be an isolating neighborhood of $Inv(N \times \Lambda, \varphi^{\Lambda}$ which admits a Lyapunov pair. Such an isolating neighborhood exists by Theorem 5.5. One easily verifies that if $\Delta \subset \Lambda$ and P is a regular index pair in M with respect to φ^{Λ} then P^{Δ} is a regular index pair in M^{Δ} with respect to φ^{Δ}. In particular if $\lambda \in \Lambda$ then P_{λ} is an index pair in M^{λ} with respect to φ^{λ}.

Let P, Q be regular index pairs in M such that $P \subset \operatorname{int} Q$. We will show that there exists a neighborhood Δ of λ_0 such that if $\kappa, \lambda \in \Delta$ then $P^{\kappa} \subset Q^{\lambda}$. Indeed, if this is not true then for some $i \in \{ 1, 2 \}$ there exist sequences $\lambda_n, \kappa_n \to \lambda_0$ and $x_n \to x \in N$ such that $x_n \in P_i^{\kappa_n} \setminus Q_i^{\lambda_n}$. It follows that $(x, \lambda_0) \in P_i \setminus \operatorname{int} Q_i$, a contradiction.

By Lemma 5.8 we can choose index pairs $P, Q, \bar{P}, \bar{Q}$ in M such that $P \subset \operatorname{int} Q$, $Q \subset \operatorname{int} \bar{P}$, $\bar{P} \subset \operatorname{int} \bar{Q}$. Choose Δ_0, a neighborhood of λ_0 such that for $\lambda \in \Delta_0$

$$P^{\lambda_0} \subset Q^{\lambda} \subset \bar{P}^{\lambda_0} \subset \bar{Q}^{\lambda}.$$

and let $\iota_1, \iota_2, \iota_3$ denote the corresponding inclusion maps. By Theorem 6.2 we have that $\mathbf{U}_{\omega}(\iota_2 \iota_1) = \mathbf{U}_{\omega}(\iota_2)\mathbf{U}_{\omega}(\iota_1)$ and $\mathbf{U}_{\omega}(\iota_3 \iota_2) = \mathbf{U}_{\omega}(\iota_3)\mathbf{U}_{\omega}(\iota_2)$ are isomorphisms. It follows that also $\mathbf{U}_{\omega}(\iota_2)$ is an isomorphism. □

As a trivial consequence of Corollary 4.7 we get the following:

Proposition 7.4 *Suppose N is a neighborhood for S and N is contractible in the phase space X. Then $h_{\omega}(S, \varphi)$ is simple.* □

If $h_{\omega}(S, \varphi)$ is simple then it is the homotopy type of the wedge (as pointed spaces) of the classical Conley index and Z and it does not contain any more information than the classical index if Z is path connected.

We conclude this section by comparing Conley indices over various base maps. The following results are easy corollaries from the corresponding results in Section 4.

Proposition 7.5 *If $\omega_0, \omega_1 : X \to Z$ are continuous, and for some neighborhood N of S, $\omega_0|_N \simeq \omega_1|_N : N \to Z$ then $h_{\omega_0}(S, \varphi) = h_{\omega_1}(S, \varphi)$.* □

Let S' be another isolated invariant set for φ.

Proposition 7.6 *If $\zeta : Z \to Z'$ is continuous, Z' is Hausdorff and $h_\omega(S,\varphi) = h_\omega(S',\varphi)$ then $h_{\zeta\circ\omega}(S,\varphi) = h_{\zeta\circ\omega}(S',\varphi)$.* □

Proposition 7.7 *If $h_\omega(S,\varphi) = h_\omega(S',\varphi)$ for some ω then the classical Conley indices $h(S,\varphi)$ and $h(S',\varphi)$ are also equal each to the other.* □

Proposition 7.8 *If $h_{\mathrm{id}_X}(S,\varphi) = h_{\mathrm{id}_X}(S',\varphi)$ then also $h_\omega(S,\varphi) = h_\omega(S',\varphi)$ for every map ω.* □

Proposition 7.9 *If $\zeta : Z \to Z'$ is a homotopy equivalence, Z' is a Hausdorff space then $h_\omega(S,\varphi) = h_\omega(S',\varphi)$ if and only if $h_{\zeta\circ\omega}(S,\varphi) = h_{\zeta\circ\omega}(S',\varphi)$.* □

8 Examples

In the following examples the term "index" means $h_\omega(S)$ for a given isolated invariant set S. We describe the index by presenting its possibly simple representant. We begin with our examples from the introduction.

Example 8.1 Consider the flow pictured in Figure 1. Let the phase space X of the flow be R^3 minus the knotted curve. Let the index pair be $P = (P_1, P_2)$, where P_1 be the cylinder (with the bases at the levels $z = 0$ and $z = 1$) minus a tubular neighborhood of the curve, and let P_2 be the punctured disc at the base in the level $z = 0$ of P_1. Put $S = \mathrm{Inv}\, P_1$. Let $Z = P_1$ and ω be the natural strong deformation retraction of X onto P_1, i.e. $\omega|_{P_1}$ is the identity and $\omega_{X\setminus P_1}$ projects the point (x,y,z) onto $(x,y,0)$ if $z \le 0$, onto $(x,y,1)$ if $z \ge 1$, and then retracts (via the natural retraction in the tubular neighborhood) all the points with $z \in [0,1]$ onto the boundary of P_1. $\mathbf{U}_\omega(P)$ consists of two copies of P_1 glued in P_2. Its fundamental group is equal to the fundamental group of the doubled knot, hence it is different from the fundamental group of Z (i.e. the group of the knot, see [Ro]), hence the index of $S = \mathrm{Inv}\, P_1$ is not trivial. In particular, it follows that any continuation of S is also nonempty – this fact cannot be directly obtained from the Ważewski Theorem.

Example 8.2 Suppose X is the punctured plane, γ_1 is a homotopically nontrivial repelling periodic orbit, and γ_2 is a homotopically trivial repelling orbit. By Proposition 7.4, the index of γ_2 is simple, i.e. has the homotopy type of the one-point wedge of the classical Conley index of γ_1 and Z. In this case the classical index is the wedge of a 1-sphere and a 2-sphere. If we let Z be the unit circle and ω be the map $z \mapsto z/|z|$, then the index of γ_1 over Z has the homotopy type of the 2-torus. (This can be seen either by homotoping Z to γ_1 and homotoping ω in a corresponding way, or by continuing γ_1 to a circle in the plane, and using an annulus as P_1.) Since γ_1 and γ_2 have different indices over Z, by Theorem 2.2, they are not related by continuation.

The next example shows that weak fiberwise homotopy type is a finer classification than ordinary homotopy type.

Example 8.3 Again, suppose X is the punctured plane, Z is the unit circle and ω is the map $z \mapsto z/|z|$, but this time the periodic orbits are attracting, with γ_1 homotopically nontrivial and γ_2 homotopically trivial. We use annuli as the isolating neighborhoods and as P_1 in the index pair. In this example, since the exit set P_2 is empty, both indices over Z have the same usual homotopy type, namely the disjoint union of two circles, one coming from P_1, the other from Z. However, the projections r_1 and r_2, for γ_1 and γ_2 respectively, map the annulus into different homotopy classes in Z. It follows from Proposition 3.5 that the resulting adjunction spaces are not weakly fiberwise homotopy equivalent, so again by Theorem 2.2, γ_1 and γ_2 are not related by continuation.

Example 8.4 If X is R^3 with the z-axis removed, Z is the unit circle in the $x-y$ plane with the obvious map $\omega(x,y,z) = (x/(x^2+y^2), y/(x^2+y^2), 0)$, then the index over Z will again distinguish periodic orbits in different homotopy classes. Let γ be a hyperbolic periodic orbit winding k times around the z-axis. Let $n = 0, 1, 2$ be the dimension of the unstable manifold of the the Poincaré map. If the Poincaré map preserves orientation on the unstable manifold, then the index is a locally trivial bundle over S^1 with the wedge of k punctured spheres S^n as the fiber. Moreover, the holonomy map transitively interchanges the spheres in the fiber and preserves their orientations. In particular, if $k = 1$ then the index is $S^n \times S^1$; if $n = 0$ then it is the disjoint union of two unit circles S^1 (one of them is treated as Z) and the projection

$S^1 \to Z$ (where S^1 denotes the other circle then Z) is given by $z \mapsto z^k$. If the Poincare map reverses the orientation of the unstable mainfold then necessarily $n = 1$. In the case $k = 1$ the index is the Klein bottle; in general it is a locally trivial bundle over S^1 with the wedge of k punctured circles S^1 as the fiber and the holonomy map transitively interchanges the circles and reverses the orientation of exactly one circle.

References

[Ba] T. Bartsch, The Conley index over a space, *Math. Z.* 209(1992), 167-177.

[Bo] K. Borsuk, *Theory of Retracts*, PWN, Warszawa 1967.

[Ch] R.C. Churchill, Isolated Invariant Sets in Compact Metric Spaces, *J. Diff. Equ.* 12(1972), 330-352.

[Co] C. Conley, *Isolated Invariant Sets and the Morse Index.* CBMS Lecture Notes **38** A.M.S. Providence, R.I. 1978.

[Coh] P. M. Cohn, The complement of a finitely generated direct summand of an abelian group, *Proc. Amer. Math. Soc.* 7 (1956), 520–521.

[E] R. Engelking, General Topology, Monografie Matematyczne, Polish Scientific Publishers, Warszawa 1977.

[J1] I. M. James, General topology over a base, in: I. M. James, E. H. Kronheimer (editors),*Aspects of Topology*, Cambridge University Press, Cambridge 1985.

[J2] I. M. James, General Topology and Homotopy Theory, Springer-Verlag, New York Berlin Heidelberg Tokyo, 1984.

[J3] I. M. James, *Fibrewise Topology*, Cambridge University Press, Cambridge 1989.

[KM] T. Kaczynski and M. Mrozek, Connected simple systems and the Conley functor, preprint.

[Ro] D. Rolfson, *Knots and Links*, Publish or Perish, Houston, 1990.

[Ry] K.P. Rybakowski, *The Homotopy Index and Partial Differential Equations*, Springer-Verlag, Berlin Heidelberg New York1987.

[Sm] J. Smoller, *Shock Waves and Reaction-Diffusion Equations*, Springer-Verlag, New York Heidelberg Berlin, 1983.

[Sw] R. M. Switzer, *Algebraic Topology – Homotopy and Homology*, Springer-Verlag, New York Heidelberg Berlin, 1975.

[W] E. A. Walker, Cancellation in direct sums of groups, *Proc. Amer. Math. Soc.* 7 (1956), 898–902.

[Wh] G. W. Whitehead, Elements of Homotopy Theory, Springer-Verlag, New York Heidelberg Berlin, 1978.